

Spin Chains in $\mathcal{N} = 2$ Superconformal Theories

From the $\mathbb{Z}_2$ Quiver to Superconformal QCD

Abhijit Gadde*, Elli Pomoni† and Leonardo Rastelli‡

*C.N. Yang Institute for Theoretical Physics,
Stony Brook University,
Stony Brook, NY 11794-3840, USA*

ABSTRACT:

In this paper we find preliminary evidence that $\mathcal{N} = 2$ superconformal QCD, the $SU(N_c)$ SYM theory with $N_f = 2N_c$ fundamental hypermultiplets, might be integrable in the large N Veneziano limit. We evaluate the one-loop dilation operator in the scalar sector of the $\mathcal{N} = 2$ superconformal quiver with $SU(N_c) \times SU(N_{\tilde{c}})$ gauge group, for $N_c \equiv N_{\tilde{c}}$. Both gauge couplings g and $\tilde{g}$ are exactly marginal. This theory interpolates between the $\mathbb{Z}_2$ orbifold of $\mathcal{N} = 4$ SYM, which corresponds to $\tilde{g} = g$, and $\mathcal{N} = 2$ superconformal QCD, which is obtained for $\tilde{g} \rightarrow 0$. The planar one-loop dilation operator takes the form of a nearest-neighbor spin-chain Hamiltonian. For superconformal QCD the spin chain is of novel form: besides the color-adjoint fields ϕ_b^a , which occupy individual sites of the chain, there are “dimers” $Q_i^a \bar{Q}_b^i$ of flavor-contracted fundamental fields, which occupy two neighboring sites. We solve the two-body scattering problem of magnon excitations and study the spectrum of bound states, for general $\tilde{g}/g$. The dimeric excitations of superconformal QCD are seen to arise smoothly for $\tilde{g} \rightarrow 0$ as the limit of bound wavefunctions of the interpolating theory. Finally we check the Yang-Baxter equation for the two-magnon S-matrix. It holds as expected at the orbifold point $\tilde{g} = g$. While violated for general $\tilde{g} \neq g$, it holds again in the limit $\tilde{g} \rightarrow 0$, hinting at one-loop integrability of planar $\mathcal{N} = 2$ superconformal QCD.

*abhijit@insti.physics.sunysb.edu

†pomoni@insti.physics.sunysb.edu

‡leonardo.rastelli@stonybrook.edu

Contents

1. Introduction	2
2. Lagrangian and Symmetries	4
2.1 $\mathcal{N} = 2$ SCQCD	4
2.2 $\mathbb{Z}_2$ orbifold of $\mathcal{N} = 4$ and interpolating family of SCFTs	6
3. One-loop Dilation Operator in the Scalar Sector	10
3.1 Hamiltonian for $\mathcal{N} = 2$ super QCD	11
3.2 Magnons in the SCQCD spin chain	13
3.3 Hamiltonian for the interpolating SCFT	15
3.4 Magnons in the interpolating spin chain	17
4. Protected Spectrum	17
4.1 Protected spectrum in $\mathcal{N} = 2$ SCQCD	18
4.2 Protected spectrum in the orbifold theory	18
4.3 Away from the orbifold point: matching with $\mathcal{N} = 2$ SCQCD	20
4.3.1 $\mathbf{SU(2)}_{\mathbf{L}}$ singlets	21
4.3.2 $\mathbf{SU(2)}_{\mathbf{L}}$ non-singlets	22
5. Two-body scattering	25
5.1 $3_L \otimes 3_R$ Sector	26
5.2 $1_L \otimes 3_R$ Sector	28
5.3 $3_L \otimes 1_R$ Sector	29
5.4 $1_L \otimes 1_R$ Sector	29
5.5 Summary	31
5.6 Left/right factorization of the two-body S-matrix	31
6. Yang-Baxter Equation	34
7. Discussion	36
A. Simplified computation of the one-loop dilation operator	39
A.1 SCQCD	39
A.2 Interpolating SCFT	41
B. The Hamiltonian for SCQCD in the Dimer Picture	42

1. Introduction

The gauge/gravity duality has given crucial insights into the dynamics of four-dimensional gauge theories. The long-standing hope is to find a precise string theory description of realistic field theories such as QCD. At present however we lack a systematic procedure to find the string dual of a given gauge theory, and all well-understood dual pairs fall into the “universality class” of the original example, the duality between $\mathcal{N} = 4$ super Yang-Mills and IIB on $AdS_5 \times S^5$. These dualities are motivated by taking the decoupling limit of brane configurations in critical string theory. Field theories in this class share a few common features, for instance: all fields are in bifundamental representations of the gauge group; the a and c conformal anomaly coefficients are equal at large N ; there is an exactly marginal coupling λ such that for λ large the dual worldsheet sigma-model is weakly coupled and the gravity approximation is valid.

To break outside the $\mathcal{N} = 4$ universality class, an important case study is $\mathcal{N} = 2$ superconformal QCD, namely the $\mathcal{N} = 2$ super Yang-Mills theory with gauge group $SU(N_c)$ and $N_f = 2N_c$ fundamental hyper multiplets. There is a large number of fundamental flavors, and $a \neq c$ at large N_c . Nevertheless the theory shares with $\mathcal{N} = 4$ SYM the crucial simplifying feature of an exactly marginal gauge coupling. In a recent paper [1] we made some progress towards the AdS dual of $\mathcal{N} = 2$ SCQCD. We attacked the problem from two fronts: from the bottom-up, we performed a systematic analysis of the protected spectrum using superconformal representation theory; from the top-down, we considered the decoupling limit of known brane constructions in string theory. We concluded that the string dual is a sub-critical string background with seven geometric dimensions, containing both AdS_5 and an S^1 factor. In this paper we take the next step of the bottom-up (=field theory) analysis, by evaluating the one-loop dilation operator in the scalar sector of the theory.

Perturbative calculations of anomalous dimensions have given important clues into the nature of $\mathcal{N} = 4$ SYM. They gave the first hint for integrability of the planar theory: the one-loop dilation operator in the scalar sector is the Hamiltonian of the integrable $SO(6)$ spin chain [2] – a result later generalized to the full theory and to higher loops, using the formalism of the asymptotic Bethe ansatz (see *e.g.* [3, 4, 5, 6, 7] for a very incomplete list of references.) Remarkably, the asymptotic S-matrix of magnon excitations in the field theory spin chain can be exactly matched with the analogous S-matrix for the dual string sigma-model. Thus perturbative calculations open a window into the structure of the dual string theory.¹ It is natural to attempt the same strategy for $\mathcal{N} = 2$ SCQCD. The theory admits a large N expansion in the Veneziano sense [9]: the number of colors N_c and the number of fundamental flavors N_f are both sent to infinity keeping fixed their ratio ($N_f/N_c \equiv 2$ in our case) and the combination $\lambda = g_{YM}^2 N_c$. We focus on the flavor-singlet sector of the theory, which is a consistent truncation since flavor singlets close under operator product expansion. Let us denote a generic color-adjoint field by ϕ_b^a , with $a, b = 1, \dots, N_c$, and a generic color-fundamental and flavor-fundamental field by Q_i^a , where $i = 1, \dots, N_f$; we are suppressing all other quantum

¹The calculation of the circular Wilson loop by localization techniques [8] is another interesting probe of the dual theory.

numbers. In the Veneziano limit, single-trace “glueball” operators, of the schematic form $\text{Tr } \phi^\ell$, are *not* closed under the action of the dilation operator – this is a major difference with respect to the standard ’t Hooft limit of large N_c with N_f fixed [10]. Rather, glueball operators mix at order one (in the large N counting) with flavor-singlet meson operators of the form $\sum_i \bar{Q}^i \phi^k Q_i$. The simplest example is the mixing of $\text{Tr}(\phi \bar{\phi})$ with the singlet meson $\sum_i \bar{Q}^i Q_i$, which occurs at one-loop in planar perturbation theory (order $O(\lambda)$). The basic “elementary” operators are thus what we call *generalized single-trace* operators, of the schematic form

$$\text{Tr} \left(\phi^{k_1} \mathcal{M}^{\ell_1} \phi^{k_2} \dots \phi^{k_n} \mathcal{M}^{\ell_n} \right), \quad \mathcal{M}_b^a \equiv \sum_{i=1}^{N_f} Q_i^a \bar{Q}_b^i, \quad (1.1)$$

where Tr is a color trace. We have introduced a flavor-contracted combination of a fundamental and an antifundamental field, $\mathcal{M}_b^a$, which for the purpose of the large N expansion plays the role of just another color-adjoint field. The usual large N factorization theorems apply: correlators of generalized multi-traces factorize into correlators of generalized single-traces. In particular, acting with the dilation operator on a generalized single-trace operator yields (at leading order in N) another generalized single-trace operator, so we may consistently diagonalize the dilation operator in the space of generalized single-traces. The dilation operator acting on generalized single-traces can then be interpreted, in the usual fashion, as the Hamiltonian of a closed spin chain. Just as in the ’t Hooft limit, planarity of the perturbative diagrams translates into locality of the spin chain: at one-loop the spin chain has only nearest neighbor interactions, at two two-loops there are next-to-nearest neighbors interactions, and each higher loop spreads the range interaction one site further.

More insight is gained by viewing $\mathcal{N} = 2$ SCQCD as part of an “interpolating” $\mathcal{N} = 2$ superconformal field theory (SCFT) that has a product gauge group $SU(N_c) \times SU(N_{\tilde{c}})$, with $N_{\tilde{c}} \equiv N_c$, and correspondingly two exactly marginal couplings g and $\tilde{g}$. For $\tilde{g} \rightarrow 0$ one recovers $\mathcal{N} = 2$ SCQCD *plus* a decoupled free vector multiplet, while for $\tilde{g} = g$ one finds the familiar $\mathbb{Z}_2$ orbifold of $\mathcal{N} = 4$ SYM. We have evaluated the one-loop dilation operator for the whole interpolating theory, in the sector of operators made out of scalar fields. The magnon excitations of the spin chain and their bound states undergo an interesting evolution as a function of $\kappa = \tilde{g}/g$. For $\kappa = 0$ (that is, for $\mathcal{N} = 2$ SCQCD itself), the basic asymptotic excitations of the spin chain are linear combinations of the the adjoint impurity $\bar{\phi}$ and of “dimer” impurities $\mathcal{M}_b^a$ (we refer to them as dimers since they occupy two sites of the chain). From the point of view of the interpolating theory with $\kappa > 0$, these dimeric asymptotic states of $\mathcal{N} = 2$ SCQCD are *bound states* of two elementary magnons; the bound-state wavefunction localizes in the limit $\kappa \rightarrow 0$, giving an impurity that occupies two sites.

Armed with the one-loop Hamiltonian in the scalar sector, we can easily determine the complete spectrum of one-loop protected composite operators made of scalar fields. It is instructive to follow the evolution of the protected eigenstates as a function of κ , from the orbifold point to $\mathcal{N} = 2$ SCQCD. Some of these results were quoted with no derivation in our previous paper [1], where they served as input to the analysis of the full protected spectrum,

carried out with the help of the superconformal index [11].

An important question is whether the one-loop spin chain of $\mathcal{N} = 2$ SCQCD is integrable. The spin chain for the $\mathbb{Z}_2$ orbifold of $\mathcal{N} = 4$ SYM (which by definition has $\check{g} = g$) is known to be integrable [12, 13]. We find that as we move away from the orbifold point integrability is broken, indeed for general $\kappa = \check{g}/g$ the Yang-Baxter equation for the two-magnon S-matrix does not hold. Remarkably however the Yang-Baxter equation is satisfied again in the $\mathcal{N} = 2$ SCQCD limit $\kappa \rightarrow 0$. Ordinarily a check of the Yang-Baxter equation is strong evidence in favor of integrability. In our case things are more subtle: the elementary Q excitations freeze in the limit $\kappa \rightarrow 0$ (their dispersion relation becomes constant), while some (but not all) of their dimeric bound states retain non-trivial dynamics. Nevertheless, for infinitesimal κ the elementary Q s are propagating excitations, and the Yang-Baxter equation fails only infinitesimally, so it seems plausible that one can define consistent Bethe equations by taking small κ as a regulator, to be removed at the end of the calculation.

In section 2 we review the Lagrangian and symmetries of $\mathcal{N} = 2$ SCQCD and of the interpolating superconformal field theory. In section 3.1 we evaluate the one-loop dilation operator of SCQCD (in the scalar sector), and write it as a spin-chain Hamiltonian. In section 3.2 we find the spectrum of magnon excitations of this spin chain. These calculations are repeated in sections 3.3 and 3.4 for the the interpolating SCFT. A simplified derivation of the Hamiltonians is presented in appendix A, while appendix B contains an equivalent way to write the Hamiltonian for $\mathcal{N} = 2$ SCQCD in terms of composite (dimeric) impurities. In section 4 we study the spectrum of protected operators of the interpolating theory, and follow its evolution in the limit $\kappa \rightarrow 0$. In section 5 we solve the two-magnon scattering problem and analyze the spectrum of bound states in the spin chain of the interpolating SCFT, with particular attention to the $\kappa \rightarrow 0$ limit. In section 5 we check the Yang-Baxter equation for the two-body S-matrix of the interpolating theory, finding that it holds for $\kappa = 1$ and $\kappa \rightarrow 0$. We conclude in section 6 with a brief discussion of integrability and of future directions of research.

2. Lagrangian and Symmetries

2.1 $\mathcal{N} = 2$ SCQCD

Our main interest is $\mathcal{N} = 2$ SYM theory with gauge group $SU(N_c)$ and $N_f = 2N_c$ fundamental hypermultiplets. We refer to this theory as $\mathcal{N} = 2$ superconformal QCD (SCQCD). Its global symmetry group is $U(N_f) \times SU(2)_R \times U(1)_r$, where $SU(2)_R \times U(1)_r$ is the R-symmetry subgroup of the superconformal group. We use indices $\mathcal{I}, \mathcal{J} = \pm$ for $SU(2)_R$, $i, j = 1, \dots, N_f$ for the flavor group $U(N_f)$ and $a, b = 1, \dots, N_c$ for the color group $SU(N_c)$.

Table 1 summarizes the field content and quantum numbers of the model: The Poincaré supercharges $Q_{\alpha}^{\mathcal{I}}$, $\bar{Q}_{\mathcal{I}\dot{\alpha}}$ and the conformal supercharges $\mathcal{S}_{\mathcal{I}\alpha}$, $\bar{\mathcal{S}}_{\dot{\alpha}}^{\mathcal{I}}$ are $SU(2)_R$ doublets with charges $\pm 1/2$ under $U(1)_r$. The $\mathcal{N} = 2$ vector multiplet consists of a gauge field A_{μ} , two Weyl spinors $\lambda_{\alpha}^{\mathcal{I}}$, $\mathcal{I} = \pm$, which form a doublet under $SU(2)_R$, and one complex scalar ϕ , all in the adjoint representation of $SU(N_c)$. Each $\mathcal{N} = 2$ hypermultiplet consists of an $SU(2)_R$ doublet

	$SU(N_c)$	$U(N_f)$	$SU(2)_R$	$U(1)_r$
$\mathcal{Q}_\alpha^{\mathcal{I}}$	1	1	2	$+1/2$
$\mathcal{S}_{\mathcal{I}\alpha}$	1	1	2	$-1/2$
A_μ	Adj	1	1	0
ϕ	Adj	1	1	-1
$\lambda_\alpha^{\mathcal{I}}$	Adj	1	2	$-1/2$
$Q_{\mathcal{I}}$	$\square$	$\square$	2	0
ψ_α	$\square$	$\square$	1	$+1/2$
$\tilde{\psi}_\alpha$	$\bar{\square}$	$\bar{\square}$	1	$+1/2$
$\mathcal{M}_1$	Adj + 1	1	1	0
$\mathcal{M}_3$	Adj + 1	1	3	0

Table 1: Symmetries of $\mathcal{N} = 2$ SCQCD. We show the quantum numbers of the supercharges $\mathcal{Q}^{\mathcal{I}}$, $\mathcal{S}_{\mathcal{I}}$, of the elementary components fields and of the mesonic operators $\mathcal{M}$. Conjugate objects (such as $\bar{\mathcal{Q}}_{\mathcal{I}\dot{\alpha}}$ and $\bar{\phi}$) are not written explicitly.

$Q_{\mathcal{I}}$ of complex scalars and of two Weyl spinors ψ_α and $\tilde{\psi}_\alpha$, $SU(2)_R$ singlets. It is convenient to define the flavor contracted mesonic operators

$$\mathcal{M}_{\mathcal{J}b}^{\mathcal{I}a} \equiv \frac{1}{\sqrt{2}} Q_{\mathcal{J}i}^a \bar{Q}_b^{\mathcal{I}i}, \quad (2.1)$$

which may be decomposed into the $SU(2)_R$ singlet and triplet combinations

$$\mathcal{M}_1 \equiv \mathcal{M}_{\mathcal{I}}^{\mathcal{I}} \quad \text{and} \quad \mathcal{M}_3 \equiv \mathcal{M}_{\mathcal{J}}^{\mathcal{I}} - \frac{1}{2} \mathcal{M}_{\mathcal{K}}^{\mathcal{K}} \delta_{\mathcal{J}}^{\mathcal{I}}. \quad (2.2)$$

The operators $\mathcal{M}$ decompose into the adjoint plus the singlet representations of the color group $SU(N_c)$; the singlet piece is however subleading in the large N_c limit.

The Lagrangian is

$$\mathcal{L} = \mathcal{L}_V + \mathcal{L}_H, \quad (2.3)$$

where $\mathcal{L}_V$ stands for the Lagrangian of the $\mathcal{N} = 2$ vector multiplet and the $\mathcal{L}_H$ for the Lagrangian of $\mathcal{N} = 2$ hypermultiplet. Explicitly²

$$\begin{aligned} \mathcal{L}_V = & -\text{Tr} \left[\frac{1}{4} F^{\mu\nu} F_{\mu\nu} + i \bar{\lambda}_{\mathcal{I}} \bar{\sigma}^\mu D_\mu \lambda^{\mathcal{I}} + (D^\mu \phi)(D_\mu \phi)^\dagger \right. \\ & \left. + i\sqrt{2} (g_{YM} \epsilon_{\mathcal{I}\mathcal{J}} \lambda^{\mathcal{I}} \lambda^{\mathcal{J}} \phi^\dagger - g_{YM} \epsilon^{\mathcal{I}\mathcal{J}} \bar{\lambda}_{\mathcal{I}} \bar{\lambda}_{\mathcal{J}} \phi) + \frac{g_{YM}^2}{2} [\phi, \phi^\dagger]^2 \right]. \end{aligned} \quad (2.4)$$

²In our conventions, $D_\mu \equiv \partial_\mu + ig_{YM} A_\mu$. We raise and lower $SU(2)_R$ indices with the antisymmetric symbols $\epsilon_{\mathcal{I}\mathcal{J}}$ and $\epsilon^{\mathcal{I}\mathcal{J}}$, which obey $\epsilon_{\mathcal{I}\mathcal{J}} \epsilon^{\mathcal{J}\mathcal{K}} = \delta_{\mathcal{I}}^{\mathcal{K}}$.

$$\begin{aligned}
\mathcal{L}_H = & - \left[(D^\mu \bar{Q}^{\mathcal{I}})(D_\mu Q_{\mathcal{I}}) + i \bar{\psi} \bar{\sigma}^\mu D_\mu \psi + i \tilde{\psi} \bar{\sigma}^\mu D_\mu \tilde{\psi} \right. \\
& + i\sqrt{2} (g_{YM} \epsilon^{\mathcal{I}\mathcal{J}} \bar{\psi} \bar{\lambda}_{\mathcal{I}} Q_{\mathcal{J}} - g_{YM} \epsilon_{\mathcal{I}\mathcal{J}} \bar{Q}^{\mathcal{I}} \lambda^{\mathcal{J}} \psi \\
& + g_{YM} \tilde{\psi} \lambda^{\mathcal{I}} Q_{\mathcal{I}} - g_{YM} \bar{Q}^{\mathcal{I}} \bar{\lambda}_{\mathcal{I}} \tilde{\psi} \\
& + g_{YM} \tilde{\psi} \phi \psi - g_{YM} \bar{\psi} \bar{\phi} \tilde{\psi}) \\
& \left. + g_{YM}^2 \bar{Q}_{\mathcal{I}} (\phi^\dagger \phi + \phi \phi^\dagger) Q^{\mathcal{I}} + g_{YM}^2 \mathcal{V}(Q) \right],
\end{aligned} \tag{2.5}$$

where the potential for the squarks is

$$\begin{aligned}
\mathcal{V}(Q) = & (\bar{Q}_a^{\mathcal{I}} Q_{\mathcal{I}}^a)(\bar{Q}_b^{\mathcal{J}} Q_{\mathcal{J}}^b) - \frac{1}{2} (\bar{Q}_a^{\mathcal{I}} Q_{\mathcal{J}}^a)(\bar{Q}_b^{\mathcal{J}} Q_{\mathcal{I}}^b) \\
& + \frac{1}{N_c} \left(\frac{1}{2} (\bar{Q}_a^{\mathcal{I}} Q_{\mathcal{I}}^a)(\bar{Q}_b^{\mathcal{J}} Q_{\mathcal{J}}^b) - (\bar{Q}_a^{\mathcal{I}} Q_{\mathcal{J}}^a)(\bar{Q}_b^{\mathcal{J}} Q_{\mathcal{I}}^b) \right).
\end{aligned} \tag{2.6}$$

Using the flavor contracted mesonic operator (2.1), $\mathcal{V}$ can be written more compactly as

$$\begin{aligned}
\mathcal{V} = & \text{Tr}[\mathcal{M}^{\mathcal{J}}_{\mathcal{I}} \mathcal{M}^{\mathcal{I}}_{\mathcal{J}}] - \frac{1}{2} \text{Tr}[\mathcal{M}^{\mathcal{I}}_{\mathcal{I}} \mathcal{M}^{\mathcal{J}}_{\mathcal{J}}] \\
& - \frac{1}{N_c} \text{Tr}[\mathcal{M}^{\mathcal{J}}_{\mathcal{I}}] \text{Tr}[\mathcal{M}^{\mathcal{I}}_{\mathcal{J}}] + \frac{1}{2} \frac{1}{N_c} \text{Tr}[\mathcal{M}^{\mathcal{I}}_{\mathcal{I}}] \text{Tr}[\mathcal{M}^{\mathcal{J}}_{\mathcal{J}}] \\
= & \text{Tr}[\mathcal{M}_3 \mathcal{M}_3] - \frac{1}{N_c} \text{Tr}[\mathcal{M}_3] \text{Tr}[\mathcal{M}_3].
\end{aligned}$$

2.2 $\mathbb{Z}_2$ orbifold of $\mathcal{N} = 4$ and interpolating family of SCFTs

$\mathcal{N} = 2$ SCQCD can be viewed as a limit of a family of superconformal theories; in the opposite limit the family reduces to a $\mathbb{Z}_2$ orbifold of $\mathcal{N} = 4$ SYM. In this subsection we first describe the orbifold theory and then its connection to $\mathcal{N} = 2$ SCQCD.

As familiar, the field content of $\mathcal{N} = 4$ SYM comprises the gauge field A_μ , four Weyl fermions λ_α^A and six real scalars X_{AB} , where $A, B = 1, \dots, 4$ are indices of the $SU(4)_R$ R-symmetry group. Under $SU(4)_R$, the fermions are in the **4** representation, while the scalars are in **6** (antisymmetric self-dual) and obey the reality condition³

$$X_{AB}^\dagger = \frac{1}{2} \epsilon^{ABCD} X_{CD}. \tag{2.7}$$

We may parametrize X_{AB} in terms of six real scalars X_k , $k = 4, \dots, 9$,

$$X_{AB} = \frac{1}{\sqrt{2}} \left(\begin{array}{cc|cc} 0 & X_4 + iX_5 & X_7 + iX_6 & X_8 + iX_9 \\ -X_4 - iX_5 & 0 & X_8 - iX_9 & -X_7 + iX_6 \\ \hline -X_7 - iX_6 & -X_8 + iX_9 & 0 & X_4 - iX_5 \\ -X_8 - iX_9 & X_7 - iX_6 & -X_4 + iX_5 & 0 \end{array} \right) \tag{2.8}$$

³The $\dagger$ indicates hermitian conjugation of the matrix in color space. We choose hermitian generators for the color group.

Next, we pick an $SU(2)_L \times SU(2)_R \times U(1)_r$ subgroup of $SU(4)_R$,

$$\begin{array}{c} 1 + \\ 2 - \\ 3 \hat{+} \\ 4 \hat{-} \end{array} \left(\begin{array}{c|c} SU(2)_R \times U(1)_r & \\ \hline & SU(2)_L \times U(1)_r^* \end{array} \right). \quad (2.9)$$

We use indices $\mathcal{I}, \mathcal{J} = \pm$ for $SU(2)_R$ (corresponding to $A, B = 1, 2$) and indices $\hat{\mathcal{I}}, \hat{\mathcal{J}} = \hat{\pm}$ for $SU(2)_L$ (corresponding to $A, B = 3, 4$). To make more manifest their transformation properties, the scalars are rewritten as the $SU(2)_L \times SU(2)_R$ singlet Z (with charge -1 under $U(1)_r$) and as the bifundamental $\mathcal{X}_{\mathcal{I}\hat{\mathcal{I}}}$ (neutral under $U(1)_r$),

$$Z \equiv \frac{X_4 + iX_5}{\sqrt{2}}, \quad \mathcal{X}_{\mathcal{I}\hat{\mathcal{I}}} \equiv \frac{1}{\sqrt{2}} \begin{pmatrix} X_7 + iX_6 & X_8 + iX_9 \\ X_8 - iX_9 & -X_7 + iX_6 \end{pmatrix}. \quad (2.10)$$

Note the reality condition $\mathcal{X}_{\mathcal{I}\hat{\mathcal{I}}}^\dagger = -\epsilon^{\mathcal{I}\mathcal{J}} \epsilon^{\hat{\mathcal{I}}\hat{\mathcal{J}}} \mathcal{X}_{\mathcal{J}\hat{\mathcal{J}}}$. Geometrically, $SU(2)_L \times SU(2)_R \cong SO(4)$ is the group of 6789 rotations and $U(1)_R \cong SO(2)$ the group of 45 rotations. Diagonal $SU(2)$ transformations $\mathcal{X} \rightarrow U\mathcal{X}U^{-1}$ ($U_R = U, U_L = U^*$) preserve the trace, $\text{Tr}[\mathcal{X}] = 2iX_6$, and thus correspond to 789 rotations.

We are now ready to discuss the orbifold projection. In R-symmetry space, the orbifold group is chosen to be $\mathbb{Z}_2 \subset SU(2)_L$ with elements $\pm \mathbb{I}_{2 \times 2}$. This is the well-known quiver theory [14] obtained by placing N_c D3 branes at the A_1 singularity $\mathbb{R}^2 \times \mathbb{R}^4/\mathbb{Z}_2$, with $(X_6, X_7, X_8, X_9) \rightarrow \pm(X_6, X_7, X_8, X_9)$ and X_4 and X_5 invariant. Supersymmetry is broken to $\mathcal{N} = 2$, since the supercharges with $SU(2)_L$ indices are projected out. The $SU(4)_R$ symmetry is broken to $SU(2)_L \times SU(2)_R \times U(1)_r$, or more precisely to $SO(3)_L \times SU(2)_R \times U(1)_r$ since only objects with integer $SU(2)_L$ spin survive. The $SU(2)_R \times U(1)_r$ factors are the R-symmetry of the unbroken $\mathcal{N} = 2$ superconformal group, while $SO(3)_L$ is an extra global symmetry under which the unbroken supercharges are neutral.

In color space, we start with gauge group $SU(2N_c)$, and declare the non-trivial element of the orbifold to be

$$\gamma \equiv \begin{pmatrix} \mathbb{I}_{N_c \times N_c} & 0 \\ 0 & -\mathbb{I}_{N_c \times N_c} \end{pmatrix}. \quad (2.11)$$

All in all the $\mathbb{Z}_2$ action on the $\mathcal{N} = 4$ fields is

$$A_\mu \rightarrow \gamma A_\mu \gamma, \quad Z_{\mathcal{I}\mathcal{J}} \rightarrow \gamma Z_{\mathcal{I}\mathcal{J}} \gamma, \quad \lambda_{\mathcal{I}} \rightarrow \gamma \lambda_{\mathcal{I}} \gamma, \quad \mathcal{X}_{\mathcal{I}\hat{\mathcal{I}}} \rightarrow -\gamma \mathcal{X}_{\mathcal{I}\hat{\mathcal{I}}} \gamma, \quad \lambda_{\hat{\mathcal{I}}} \rightarrow -\gamma \lambda_{\hat{\mathcal{I}}} \gamma. \quad (2.12)$$

	$SU(N_c)$	$SU(N_{\tilde{c}})$	$SU(2)_R$	$SU(2)_L$	$U(1)_R$
$\mathcal{Q}_\alpha^{\mathcal{I}}$	1	1	2	1	+1/2
$\mathcal{S}_{\mathcal{I}\alpha}$	1	1	2	1	-1/2
A_μ	Adj	1	1	1	0
$\check{A}_\mu$	1	Adj	1	1	0
ϕ	Adj	1	1	1	-1
$\check{\phi}$	1	Adj	1	1	-1
$\lambda^{\mathcal{I}}$	Adj	1	2	1	-1/2
$\check{\lambda}^{\mathcal{I}}$	1	Adj	2	1	-1/2
$Q_{\mathcal{I}\hat{\mathcal{I}}}$	$\square$	$\overline{\square}$	2	2	0
$\psi_{\hat{\mathcal{I}}}$	$\square$	$\overline{\square}$	1	2	+1/2
$\check{\psi}_{\hat{\mathcal{I}}}$	$\overline{\square}$	$\square$	1	2	+1/2

Table 2: Symmetries of the $\mathbb{Z}_2$ orbifold of $\mathcal{N} = 4$ SYM and of the interpolating family of $\mathcal{N} = 2$ SCFTs.

The components that survive the projection are

$$A_\mu = \begin{pmatrix} A_{\mu b}^a & 0 \\ 0 & \check{A}_{\mu \check{b}}^{\check{a}} \end{pmatrix} \quad Z = \begin{pmatrix} \phi^a{}_b & 0 \\ 0 & \check{\phi}^{\check{a}}{}_{\check{b}} \end{pmatrix} \quad (2.13)$$

$$\lambda_{\mathcal{I}} = \begin{pmatrix} \lambda_{\mathcal{I}b}^a & 0 \\ 0 & \check{\lambda}_{\mathcal{I}\check{b}}^{\check{a}} \end{pmatrix} \quad \lambda_{\hat{\mathcal{I}}} = \begin{pmatrix} 0 & \psi_{\hat{\mathcal{I}}\check{a}}^a \\ \tilde{\psi}_{\hat{\mathcal{I}}b}^{\check{b}} & 0 \end{pmatrix} \quad (2.14)$$

$$\mathcal{X}_{\mathcal{I}\hat{\mathcal{I}}} = \begin{pmatrix} 0 & Q_{\mathcal{I}\hat{\mathcal{I}}\check{a}}^a \\ -\epsilon_{\mathcal{I}\mathcal{J}}\epsilon_{\hat{\mathcal{I}}\hat{\mathcal{J}}} \bar{Q}_{\check{b}\hat{\mathcal{J}}\mathcal{J}} & 0 \end{pmatrix}. \quad (2.15)$$

The gauge group is broken to $SU(N_c) \times SU(N_{\tilde{c}}) \times U(1)$, where the $U(1)$ factor is the *relative*⁴ $U(1)$ generated by γ (equ.(2.11)): it must be removed by hand, since its beta function is non-vanishing. The process of removing the relative $U(1)$ modifies the scalar potential by double-trace terms, which arise from the fact that the auxiliary fields (in $\mathcal{N} = 1$ superspace) are now missing the $U(1)$ component. Equivalently we can evaluate the beta function for the double-trace couplings, and tune them to their fixed point [15].

Supersymmetry organizes the component fields into the $\mathcal{N} = 2$ vector multiplets of each factor of the gauge group, $(\phi, \lambda_{\mathcal{I}}, A_\mu)$ and $(\check{\phi}, \check{\lambda}_{\mathcal{I}}, \check{A}_\mu)$, and into two bifundamental hypermultiplets, $(Q_{\mathcal{I},\hat{\mathcal{I}}}, \psi_{\hat{\mathcal{I}}}, \tilde{\psi}_{\hat{\mathcal{I}}})$ and $(Q_{\mathcal{I},\check{\mathcal{I}}}, \psi_{\check{\mathcal{I}}}, \tilde{\psi}_{\check{\mathcal{I}}})$. Table 2 summarizes the field content and quantum numbers of the orbifold theory.

The two gauge-couplings g_{YM} and $\check{g}_{YM}$ can be independently varied while preserving $\mathcal{N} = 2$ superconformal invariance, thus defining a two-parameter family of $\mathcal{N} = 2$ SCFTs. Some care is needed in adjusting the Yukawa and scalar potential terms so that $\mathcal{N} = 2$

⁴Had we started with $U(2N_c)$ group, we would also have an extra *diagonal* $U(1)$, which would completely decouple since no fields are charged under it.

supersymmetry is preserved. We find

$$\begin{aligned}\mathcal{L}_{Yukawa}(g_{YM}, \check{g}_{YM}) = & i\sqrt{2}\text{Tr}[-g_{YM}\epsilon^{\mathcal{I}\mathcal{J}}\bar{\lambda}_{\mathcal{I}}\bar{\lambda}_{\mathcal{J}}\phi - \check{g}_{YM}\epsilon^{\mathcal{I}\mathcal{J}}\bar{\check{\lambda}}_{\mathcal{I}}\bar{\check{\lambda}}_{\mathcal{J}}\check{\phi} \\ & + g_{YM}\epsilon^{\hat{\mathcal{I}}\hat{\mathcal{J}}}\tilde{\psi}_{\hat{\mathcal{I}}}\phi\psi_{\hat{\mathcal{J}}} + \check{g}_{YM}\epsilon^{\hat{\mathcal{I}}\hat{\mathcal{J}}}\psi_{\hat{\mathcal{J}}}\check{\phi}\tilde{\psi}_{\hat{\mathcal{I}}} \\ & + g_{YM}\epsilon^{\hat{\mathcal{I}}\hat{\mathcal{J}}}\tilde{\psi}_{\hat{\mathcal{J}}}\lambda^{\mathcal{I}}Q_{\mathcal{I}\hat{\mathcal{I}}} + \check{g}_{YM}\epsilon^{\hat{\mathcal{I}}\hat{\mathcal{J}}}\tilde{\psi}_{\hat{\mathcal{I}}}\lambda^{\mathcal{I}}Q_{\mathcal{I}\hat{\mathcal{I}}} \\ & - g_{YM}\epsilon_{\mathcal{I}\mathcal{J}}\bar{Q}^{\hat{\mathcal{J}}\mathcal{I}}\lambda^{\mathcal{J}}\psi_{\hat{\mathcal{J}}} - \check{g}_{YM}\epsilon_{\mathcal{I}\mathcal{J}}\psi_{\hat{\mathcal{J}}}\bar{\check{\lambda}}^{\mathcal{I}}\bar{Q}^{\hat{\mathcal{J}}\mathcal{I}}] + h.c.\end{aligned}\quad (2.16)$$

$$\begin{aligned}\mathcal{V}(g_{YM}, \check{g}_{YM}) = & g_{YM}^2\text{Tr}[\frac{1}{2}[\bar{\phi}, \phi]^2 + \mathcal{M}_{\mathcal{I}}^{\mathcal{I}}(\phi\bar{\phi} + \bar{\phi}\phi) + \mathcal{M}_{\mathcal{I}}^{\mathcal{J}}\mathcal{M}_{\mathcal{J}}^{\mathcal{I}} - \frac{1}{2}\mathcal{M}_{\mathcal{I}}^{\mathcal{I}}\mathcal{M}_{\mathcal{J}}^{\mathcal{J}}] \\ & + \check{g}_{YM}^2\text{Tr}[\frac{1}{2}[\bar{\check{\phi}}, \check{\phi}]^2 + \check{\mathcal{M}}_{\mathcal{I}}^{\mathcal{I}}(\check{\phi}\bar{\check{\phi}} + \bar{\check{\phi}}\check{\phi}) + \check{\mathcal{M}}_{\mathcal{I}}^{\mathcal{J}}\check{\mathcal{M}}_{\mathcal{J}}^{\mathcal{I}} - \frac{1}{2}\check{\mathcal{M}}_{\mathcal{I}}^{\mathcal{I}}\check{\mathcal{M}}_{\mathcal{J}}^{\mathcal{J}}] \\ & + g_{YM}\check{g}_{YM}\text{Tr}[-2Q_{\mathcal{I}\hat{\mathcal{I}}}\check{\phi}\bar{Q}^{\hat{\mathcal{I}}\mathcal{I}}\bar{\phi} + h.c.] - \frac{1}{N_c}\mathcal{V}_{d.t.},\end{aligned}\quad (2.17)$$

where the mesonic operators $\mathcal{M}$ are defined as⁵

$$\mathcal{M}_{\mathcal{J}}^{\mathcal{I}a} \equiv \frac{1}{\sqrt{2}}Q_{\mathcal{J}\hat{\mathcal{J}}\check{a}}^a\bar{Q}^{\hat{\mathcal{J}}\mathcal{I}a}, \quad \check{\mathcal{M}}_{\mathcal{J}\check{b}}^{\mathcal{I}\check{a}} \equiv \frac{1}{\sqrt{2}}\bar{Q}^{\hat{\mathcal{J}}\mathcal{I}\check{a}}Q_{\mathcal{J}\hat{\mathcal{J}}\check{b}}^a, \quad (2.18)$$

and the double-trace terms in the potential are

$$\begin{aligned}\mathcal{V}_{d.t.} = & g_{YM}^2(\text{Tr}[\mathcal{M}_{\mathcal{I}}^{\mathcal{J}}]\text{Tr}[\mathcal{M}_{\mathcal{J}}^{\mathcal{I}}] - \frac{1}{2}\text{Tr}[\mathcal{M}_{\mathcal{I}}^{\mathcal{I}}]\text{Tr}[\mathcal{M}_{\mathcal{J}}^{\mathcal{J}}]) \\ & + \check{g}_{YM}^2(\text{Tr}[\check{\mathcal{M}}_{\mathcal{I}}^{\mathcal{J}}]\text{Tr}[\check{\mathcal{M}}_{\mathcal{J}}^{\mathcal{I}}] - \frac{1}{2}\text{Tr}[\check{\mathcal{M}}_{\mathcal{I}}^{\mathcal{I}}]\text{Tr}[\check{\mathcal{M}}_{\mathcal{J}}^{\mathcal{J}}]) \\ = & (g_{YM}^2 + \check{g}_{YM}^2)(\text{Tr}[\mathcal{M}_{\mathcal{I}}^{\mathcal{J}}]\text{Tr}[\mathcal{M}_{\mathcal{J}}^{\mathcal{I}}] - \frac{1}{2}\text{Tr}[\mathcal{M}_{\mathcal{I}}^{\mathcal{I}}]\text{Tr}[\mathcal{M}_{\mathcal{J}}^{\mathcal{J}}]).\end{aligned}\quad (2.19)$$

The $SU(2)_L$ symmetry is present for all values of the couplings (and so is the $SU(2)_R \times U(1)_r$ R-symmetry, of course). At the orbifold point $g_{YM} = \check{g}_{YM}$ there is an extra $\mathbb{Z}_2$ symmetry (the quantum symmetry of the orbifold) acting as

$$\phi \leftrightarrow \check{\phi}, \quad \lambda_{\mathcal{I}} \leftrightarrow \check{\lambda}_{\mathcal{I}}, \quad A_{\mu} \leftrightarrow \check{A}_{\mu}, \quad \psi_{\hat{\mathcal{I}}} \leftrightarrow \tilde{\psi}_{\hat{\mathcal{I}}}, \quad Q_{\mathcal{I}\hat{\mathcal{I}}} \leftrightarrow -\epsilon_{\mathcal{I}\mathcal{J}}\epsilon_{\hat{\mathcal{I}}\hat{\mathcal{J}}}\bar{Q}^{\mathcal{J}\hat{\mathcal{J}}}. \quad (2.20)$$

Setting $\check{g}_{YM} = 0$, the second vector multiplet $(\check{\phi}, \check{\lambda}_{\mathcal{I}}, \check{A}_{\mu})$ becomes free and completely decouples from the rest of theory, which happens to coincide with $\mathcal{N} = 2$ SCQCD (indeed the field content is the same and $\mathcal{N} = 2$ susy does the rest). The $SU(N_{\check{c}})$ symmetry can now be interpreted as a global flavor symmetry. In fact there is a symmetry enhancement $SU(N_{\check{c}}) \times SU(2)_L \rightarrow U(N_f = 2N_c)$: one sees in (2.16, 2.17) that for $\check{g}_{YM} = 0$ the $SU(N_{\check{c}})$ index $\check{a}$ and the $SU(2)_L$ index $\hat{\mathcal{I}}$ can be combined into a single flavor index $i \equiv (\check{a}, \hat{\mathcal{I}}) = 1, \dots, 2N_c$.

In the rest of the paper, unless otherwise stated, we will work in the large $N_c \equiv N_{\check{c}}$ limit,

⁵Note that $\text{Tr}[\mathcal{M}_{\mathcal{I}}^{\mathcal{J}}] = \text{Tr}[\check{\mathcal{M}}_{\mathcal{I}}^{\mathcal{J}}]$.

keeping fixed the ‘t Hooft couplings

$$\lambda \equiv g_{YM}^2 N_c \equiv 8\pi^2 g^2, \quad \check{\lambda} \equiv \check{g}_{YM}^2 N_{\check{c}} \equiv 8\pi^2 \check{g}^2. \quad (2.21)$$

We will refer to the theory with arbitrary g and $\check{g}$ as the “interpolating SCFT”, thinking of keeping g fixed as we vary $\check{g}$ from $\check{g} = g$ (orbifold theory) to $\check{g} = 0$ ($\mathcal{N} = 2$ SCQCD $\oplus$ extra $N_{\check{c}}^2 - 1$ free vector multiplets).

3. One-loop Dilation Operator in the Scalar Sector

At large $N_c \sim N_f$, the dilation operator of $\mathcal{N} = 2$ SCQCD can be diagonalized in the sector of generalized single-trace operators, of the form (1.1), indeed the mixing with generalized multi-traces is subleading. Motivated by the success of the analogous calculation in $\mathcal{N} = 4$ SYM [2], we have evaluated the one-loop dilation operator on generalized single-trace operators made out of scalar fields. An example of such an operator is

$$\text{Tr}[\bar{\phi}\phi\phi Q_{\mathcal{I}}\bar{Q}^{\mathcal{J}}\bar{\phi}] = \bar{\phi}^a_b \phi^b_c \phi^c_d Q_{\mathcal{I}}^d \bar{Q}^{\mathcal{J}i}_e \bar{\phi}^e_a, \quad a, b, c, d, e = 1, \dots, N_c, \quad i = 1, \dots, N_f. \quad (3.1)$$

Since the color or flavor indices of consecutive elementary fields are contracted, we can assign each field to a definite “lattice site”⁶ and think of a generalized single-trace operator as a state in a periodic spin chain. In the scalar sector, the state space V_l at each lattice site is six-dimensional, spanned by $\{\phi, \bar{\phi}, Q_{\mathcal{I}}, \bar{Q}^{\mathcal{J}}\}$. However the index structure of the fields imposes restrictions on the total space $\otimes_{l=1}^L V_l$: not all states in the tensor product are allowed. Indeed a Q at site l must always be followed by a $\bar{Q}$ at site $l + 1$, and viceversa a $\bar{Q}$ must always be preceded by a Q . Equivalently, as in appendix B, we may use instead the color-adjoint objects $\phi, \bar{\phi}, \mathcal{M}_1$ and $\mathcal{M}_3$ (recall the definitions (2.2)), where the $\mathcal{M}$ ’s are viewed as “dimers” occupying two sites of the chain.

As usual, we may interpret the perturbative dilation operator as the Hamiltonian of the spin chain. It is convenient to factor out the overall coupling from the definition of the Hamiltonian H ,

$$\Gamma^{(1)} \equiv g^2 H, \quad g^2 \equiv \frac{\lambda}{8\pi^2}, \quad \lambda \equiv g_{YM}^2 N_c, \quad (3.2)$$

where $\Gamma^{(1)}$ is the one-loop anomalous dimension matrix. By a simple extension of the usual arguments, the Veneziano double-line notation (see figure 6 for an example) makes it clear that for large $N_c \sim N_f$ (with λ fixed) the dominant contribution comes from planar diagrams. Planarity implies that the one-loop Hamiltonian is of nearest-neighbor type, $H = \sum_{l=1}^L H_{kk+1}$ (with $k \equiv k + L$), where $H_{k,k+1} : V_k \otimes V_{k+1} \rightarrow V_k \otimes V_{k+1}$. The two-loop correction is next-to-nearest-neighbor and so on. In section 3.1 we present our results for the one-loop Hamiltonian of the spin chain for SCQCD. We then derive (section 3.2) the one-particle “magnon” excitations of the infinite chain above the BPS vacuum $\dots \phi\phi\phi\dots$. The one-particle

⁶Up to cyclic re-ordering of course, under which the trace is invariant.




Figure 1: Various types of Feynman diagrams that contribute, at one loop, to anomalous dimension. The first diagram is the self-energy contribution. The second diagram represents the gluon exchange contribution whereas the third one stands for the quartic interaction between the fields. The first and the second diagrams are proportional to the identity in the R symmetry space while the third one carries a nontrivial R symmetry index structure.

eigenstates are interesting admixtures of the adjoint $\bar{\phi}$ impurity and of the “dimeric” $Q\bar{Q}$ impurities.

The generalization to the full interpolating SCFT is straightforward and is carried out in sections 3.3 and 3.4. The structure of this more general spin chain is in a sense more conventional, and it is somewhat reminiscent of the spin chain [16, 17, 18, 19] for the ABJM [20] and ABJ [21] theories.⁷

There are two types of color indices, for the two gauge groups $SU(N_c)$ and $SU(N_{\bar{c}})$, with adjoint fields ϕ_b^a and $\check{\phi}_{\bar{b}}^{\bar{a}}$ carrying two indices of the same type, and bifundamental fields Q_b^a and $\bar{Q}_{\bar{b}}^{\bar{a}}$ carrying two indices of opposite type. Of course one must contract neighboring indices of the same type. Now a Q and a $\bar{Q}$ need not be adjacent since they can be separated for $\check{\phi}$ fields. The infinite chain admits two BPS vacua, the state with all ϕ s and the state with all $\check{\phi}$ s. The magnons are momentum eigenstates containing a single Q or $\bar{Q}$ impurity, separating one BPS vacuum on the left from the other vacuum on the right. We will see in section 5 how the “dimeric” $Q\bar{Q}$ impurities of the SCQCD chain arise in the limit $\check{g} \rightarrow 0$ from the localization of the bound state wavefunctions of the interpolating chain.

3.1 Hamiltonian for $\mathcal{N} = 2$ super QCD

We have determined the one-loop dilation operator in the scalar sector by explicit evaluation of the divergent part of all the relevant Feynman diagrams, which can be classified as self energy diagrams, gluon interaction diagrams and quartic vertex diagrams and are schematically shown in figure 1. The calculation is straightforward and its details will not be reproduced here. In appendix A we present a shortcut derivation that bypasses the explicit evaluation of the

⁷An important difference is that our spin chain has an exact parity symmetry, whereas the spin chain of the ABJ theory is expected to violate parity at sufficiently high perturbative order (though somewhat surprisingly the ABJ planar theory appears to be parity invariant to low perturbative order [19, 22, 23].)

self-energy and gluon exchange diagrams, whose contribution can be fixed by requiring the vanishing of the anomalous dimension of certain protected operators.

As we are at it, we may as well consider the case of arbitrary N_f , though we are ultimately interested in the conformal case $N_f = 2N_c$. In the non-conformal case, it is more useful to normalize the fields so that the Lagrangian has an overall factor of $1/g_{YM}^2$ in front [24]. This different normalization affects the anomalous dimension of composite operators for $N_f \neq 2N_c$, which acquire an extra contribution due to the beta function, but it is of course immaterial for $N_f = 2N_c$. It is in this normalization that the chiral operator $\text{Tr}\phi^\ell$ has vanishing anomalous dimension for all N_f .

We find⁸

$$H_{k,k+1} = \quad (3.3)$$

$$\begin{array}{c} \phi^{\mathbf{p}}\phi^{\mathbf{q}} \quad Q_{\mathcal{I}}\bar{Q}^{\mathcal{J}} \quad \bar{Q}^{\mathcal{K}}Q_{\mathcal{L}} \quad \bar{Q}^{\mathcal{I}}\phi^{\mathbf{p}} \quad \phi^{\mathbf{p}}Q_{\mathcal{I}} \\ \left(\begin{array}{ccccc} \phi_{\mathbf{p}'}\phi_{\mathbf{q}'} & 2\delta_{\mathbf{p}'}^{\mathbf{p}}\delta_{\mathbf{q}'}^{\mathbf{q}} + g^{\mathbf{p}\mathbf{q}}g_{\mathbf{p}'\mathbf{q}'} - 2\delta_{\mathbf{q}'}^{\mathbf{p}}\delta_{\mathbf{p}'}^{\mathbf{q}} & \sqrt{\frac{N_f}{N_c}}g_{\mathbf{p}'\mathbf{q}'}\delta_{\mathcal{I}}^{\mathcal{J}} & 0 & 0 \\ \bar{Q}^{\mathcal{I}'}Q_{\mathcal{J}'} & \sqrt{\frac{N_f}{N_c}}g^{\mathbf{p}\mathbf{q}}\delta_{\mathcal{J}'}^{\mathcal{I}'} & (2\delta_{\mathcal{I}}^{\mathcal{J}'}\delta_{\mathcal{J}'}^{\mathcal{I}} - \delta_{\mathcal{I}}^{\mathcal{J}}\delta_{\mathcal{J}'}^{\mathcal{I}'})\frac{N_f}{N_c} & 0 & 0 \\ Q_{\mathcal{K}'}\bar{Q}^{\mathcal{L}'} & 0 & 0 & 2\delta_{\mathcal{L}}^{\mathcal{K}'}\delta_{\mathcal{K}'}^{\mathcal{L}'} & 0 \\ Q_{\mathcal{I}'}\phi_{\mathbf{p}'} & 0 & 0 & -\frac{1}{2}(1+\xi)\delta_{\mathcal{K}'}^{\mathcal{K}}\delta_{\mathcal{L}}^{\mathcal{L}'} & \frac{1}{4}(7-\xi)\delta_{\mathcal{I}}^{\mathcal{I}'}\delta_{\mathbf{p}'}^{\mathbf{p}} \\ \phi_{\mathbf{p}'}\bar{Q}^{\mathcal{I}'} & 0 & 0 & 0 & \frac{1}{4}(7-\xi)\delta_{\mathcal{I}}^{\mathcal{I}'}\delta_{\mathbf{p}'}^{\mathbf{p}} \end{array} \right) \end{array}$$

The indices $\mathbf{p}, \mathbf{q} = \pm$ label the $U(1)_r$ charges of ϕ and $\bar{\phi}$, in other terms we have defined $\phi^- \equiv \phi$, $\phi^+ \equiv \bar{\phi}$, and $g_{\mathbf{p}\mathbf{q}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. The parameter ξ is the gauge parameter that appears in the gluon propagator as $\frac{1}{k^2}(g_{\mu\nu} - (1-\xi)\frac{k_\mu k_\nu}{k^2})$. Although the form of nearest-neighbor Hamiltonian depends on gauge choice ξ , it is easy to check that ξ dependence drops when H acts on a closed chain. In the following we will set $\xi = -1$.⁹

⁸The spin chain with this nearest-neighbor Hamiltonian reproduces the one-loop anomalous dimension of all operators with $L > 2$, where L is the number of sites. The $L = 2$ case is special: the double-trace terms in the scalar potential, which give subleading contributions (at large N) for $L > 2$, become important for $L = 2$ and must be added separately. This special case plays a role in the protection of $\text{Tr}\mathcal{M}_3$, see section 4.

⁹This choice corresponds to setting to zero the self-energy of Q and $\bar{Q}$. Then our Hamiltonian can also be used as is to calculate the anomalous dimension of operators with open *flavor* indices, of the schematic form $\bar{Q}^i \dots Q_j$. For $\xi \neq -1$ there are extra contributions from the self-energy of the Q^i and $\bar{Q}_j$ at the edge of the chain.

We may rewrite H_{kk+1} more concisely (we have set $\xi = -1$) as

$$H_{k,k+1} = \begin{matrix} & \phi\phi & Q\bar{Q} & \bar{Q}Q & \bar{Q}\phi & \phi Q \\ \begin{matrix} \phi\phi \\ \bar{Q}Q \\ Q\bar{Q} \\ Q\phi \\ \phi\bar{Q} \end{matrix} & \begin{pmatrix} 2\mathbb{I} + \mathbb{K} - 2\mathbb{P} & \sqrt{\frac{N_f}{N_c}} & 0 & 0 & 0 \\ \sqrt{\frac{N_f}{N_c}} & (2\mathbb{I} - \mathbb{K})\frac{N_f}{N_c} & 0 & 0 & 0 \\ 0 & 0 & 2\mathbb{K} & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 2 \end{pmatrix} \end{matrix} \quad (3.4)$$

The symbols $\mathbb{I}, \mathbb{P}$ and $\mathbb{K}$ for identity, permutation and trace operators respectively. Their position in the matrix specifies the space in which they act. For example, the operator $\mathbb{P}$ that appears in the matrix element of $\langle \phi_{\mathbf{p}'} \phi_{\mathbf{q}'} | \phi^{\mathbf{p}} \phi^{\mathbf{q}} \rangle$ is $\delta_{\mathbf{q}'}^{\mathbf{p}} \delta_{\mathbf{p}'}^{\mathbf{q}}$, the operator $\mathbb{K}$ that appears in the matrix element $\langle \bar{Q}^{T'} Q_{\mathcal{J}'} | Q_{\mathcal{I}} \bar{Q}^{\mathcal{J}} \rangle$ stands for the operator $\delta_{\mathcal{J}'}^{T'} \delta_{\mathcal{I}}^{\mathcal{J}}$ and so on. The entries where no symbols appear have an unambiguous index structure. In appendix B we present an equivalent form of the Hamiltonian in terms of composite (dimeric) impurities.

Although not immediately obvious from the form (3.4), the Hamiltonian of the SCQCD spin chain preserves parity, once the constraints on the states allowed by the index structure are taken into account. Parity is in fact a symmetry of the spin chain for the whole interpolating theory, the transformation rules are given below in (3.20).

For $N_f = 0$, the Hamiltonian can be consistently truncated to the space of ϕ (and $\bar{\phi}$): it reduces to $2\mathbb{I}_{\phi\phi} + \mathbb{K}_{\phi\phi} - 2\mathbb{P}_{\phi\phi}$, which is Hamiltonian of the XXZ spin chain, confirming the result found in [24] for pure $\mathcal{N} = 2$ SYM. The $N_f \neq 0$ the ϕ sector is not closed in our case due to the leading order glueball-meson mixing.

3.2 Magnons in the SCQCD spin chain

The chiral operator $\text{Tr } \phi^\ell$ and the antichiral operator $\text{Tr } \bar{\phi}^\ell$ are zero-energy eigenstates (in particular the mixing element that is responsible for $\phi\phi \rightarrow Q\bar{Q}$ is proportional to $\mathbb{K}$ in ϕ space, and thus vanishes when two neighboring ϕ fields have the same $U(1)_r$ index). They correspond to the two ferromagnetic ground states of the spin chain (all spins up or all down). We choose for definiteness the chiral vacuum $\text{Tr } \phi^\ell$. Recall that in our conventions the $U(1)_r$ charge of ϕ is $r = -1$, so the ground state obeys $\Delta + r = 0$, where Δ is the total conformal dimension. Both Q and $\bar{Q}$ have $\Delta + r = 1$, but the index structure forbids the insertion of only one of them. The simplest impurities that can be excited on the ground state are $\bar{\phi}$, $\mathcal{M}_1$ and $\mathcal{M}_3$, where the last two are “dimeric” impurities which occupy two sites (recall (2.18)). All of them have $\Delta + r = 2$, and should be viewed in this sense as double excitations, though they are the most elementary we can find in the spin chain for $\mathcal{N} = 2$ SCQCD. We will see that they can be viewed as bound states of the elementary impurities of the interpolating theory with $\tilde{g} \neq 0$. This hidden compositeness makes the scattering problem somewhat harder than usual.

In the map from the (generalized) single-trace operators to the states of the spin chain, cyclicity of the trace gives periodic boundary conditions on the chain, along with the constraint that the total momentum of all the impurities in the spin be zero. As usual, it is convenient to first consider the chain to be infinite, and impose later the zero-momentum constraint on multi-impurity states. We now proceed to diagonalize the Hamiltonian on the space of states containing a single impurity (which in the present context means a single $\bar{\phi}$ or $\mathcal{M}_1$ or $\mathcal{M}_3$). The action of H on single impurities in position space is

$$H[\bar{\phi}(x)] = 6\bar{\phi}(x) - \bar{\phi}(x+1) - \bar{\phi}(x-1) \quad (3.5)$$

$$+ \sqrt{\frac{2N_f}{N_c}} \mathcal{M}_1(x) + \sqrt{\frac{2N_f}{N_c}} \mathcal{M}_1(x-1) \quad (3.6)$$

$$H[\mathcal{M}_1(x)] = 4\mathcal{M}_1(x) + \sqrt{\frac{2N_f}{N_c}} \bar{\phi}(x) + \sqrt{\frac{2N_f}{N_c}} \bar{\phi}(x+1) \\ H[\mathcal{M}_3(x)] = 8\mathcal{M}_3(x), \quad (3.7)$$

where the coordinate x denotes the site of the impurity on the chain; for the dimeric impurities $\mathcal{M}_1$ and $\mathcal{M}_3$ we use the coordinate of the first site. To diagonalize the Hamiltonian on the $\bar{\phi}/\mathcal{M}_1$ sector, we go to momentum space,

$$\bar{\phi}(p) \equiv \sum_x \bar{\phi}(x) e^{ipx}, \quad \mathcal{M}_1(p) \equiv \sum_x \mathcal{M}_1(x) e^{ipx} \quad (3.8)$$

$$H \begin{pmatrix} \bar{\phi}(p) \\ \mathcal{M}_1 \end{pmatrix} = \begin{pmatrix} 6 - e^{ip} - e^{-ip} & (1 + e^{-ip}) \sqrt{\frac{2N_f}{N_c}} \\ (1 + e^{ip}) \sqrt{\frac{2N_f}{N_c}} & 4 \end{pmatrix} \begin{pmatrix} \bar{\phi}(p) \\ \mathcal{M}_1 \end{pmatrix}. \quad (3.9)$$

The expressions for the eigenvalues and eigenvectors are not very illuminating for generic values of the ratio N_f/N_c . For the conformal case of $N_f = 2N_c$, however, they simplify. The eigenstates for $N_f = 2N_c$ are

$$T(p) \equiv -\frac{1}{2}(1 + e^{-ip})\bar{\phi}(p) + \mathcal{M}_1(p) = \sum_x e^{ipx} \left[-\frac{1}{2}(\bar{\phi}(x) + \bar{\phi}(x+1)) + \mathcal{M}_1(x) \right] \quad (3.10)$$

$$\tilde{T}(p) \equiv \bar{\phi}(p) + \frac{1}{2}(1 + e^{ip})\mathcal{M}_1(p) = \sum_x e^{ipx} \left[\bar{\phi}(x) + \frac{1}{2}(\mathcal{M}_1(x) + \mathcal{M}_1(x-1)) \right], \quad (3.11)$$

with eigenvalues

$$HT(p) = 4 \sin^2\left(\frac{p}{2}\right) T(p) \quad (3.12)$$

$$H\tilde{T}(p) = 8\tilde{T}(p). \quad (3.13)$$

Interestingly, precisely at the conformal point $N_f = 2N_c$ the magnon excitation $T(p)$ becomes gapless: in general the gap of $T(p)$ is $4 - 2\sqrt{2N_f/N_c}$. From now on we will only consider the

superconformal case and set $N_f \equiv 2N_c$. Besides $T(p)$ and $\tilde{T}(p)$, we have of course also the $\mathcal{M}_3$ momentum eigenstate,

$$\mathcal{M}_3(p) \equiv \sum_x \mathcal{M}_3(x) e^{ipx}, \quad (3.14)$$

which has the same momentum-independent energy as $\tilde{T}(p)$,

$$H\mathcal{M}_3(p) = 8\mathcal{M}_3(p). \quad (3.15)$$

3.3 Hamiltonian for the interpolating SCFT

We have generalized the calculation of the one-loop dilation operator to the full interpolating family of $\mathcal{N} = 2$ SCFTs, in the scalar sector. We find

$$\begin{aligned} H = & \begin{array}{cc} \phi^{\mathfrak{p}}\phi^{\mathfrak{q}} & Q_{I\hat{I}}\bar{Q}^{\hat{\mathcal{I}}\mathcal{J}} \\ \phi_{\mathfrak{p}'}\phi_{\mathfrak{q}'} & \delta_{\hat{\mathcal{I}}}^{\mathcal{J}}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}}g_{\mathfrak{p}'\mathfrak{q}'} \end{array} \left(\begin{array}{cc} (2\delta_{\mathfrak{p}'}^{\mathfrak{p}}\delta_{\mathfrak{q}'}^{\mathfrak{q}} + g^{\mathfrak{p}\mathfrak{q}}g_{\mathfrak{p}'\mathfrak{q}'} - 2\delta_{\mathfrak{q}'}^{\mathfrak{p}}\delta_{\mathfrak{p}'}^{\mathfrak{q}}) & \\ \delta_{\mathcal{J}'}^{\mathcal{I}'}\delta_{\hat{\mathcal{J}}'}^{\hat{\mathcal{I}}'}g^{\mathfrak{p}\mathfrak{q}} & (2\delta_{\hat{\mathcal{I}}}^{\mathcal{I}'}\delta_{\mathcal{J}'}^{\mathcal{J}} - \delta_{\hat{\mathcal{I}}}^{\mathcal{J}}\delta_{\mathcal{J}'}^{\mathcal{I}'})\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}}\delta_{\hat{\mathcal{J}}'}^{\hat{\mathcal{I}}'} + 2\kappa^2\delta_{\hat{\mathcal{I}}}^{\mathcal{J}}\delta_{\mathcal{J}'}^{\mathcal{I}'}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{I}}'}\delta_{\hat{\mathcal{J}}'}^{\hat{\mathcal{J}}} \end{array} \right) \\ & \oplus \begin{array}{cc} \check{\phi}^{\mathfrak{p}}\check{\phi}^{\mathfrak{q}} & \bar{Q}^{\hat{\mathcal{I}}\mathcal{J}}Q_{I\hat{I}} \\ \check{\phi}_{\mathfrak{p}'}\check{\phi}_{\mathfrak{q}'} & \kappa^2\delta_{\hat{\mathcal{I}}}^{\mathcal{J}}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}}g_{\mathfrak{p}'\mathfrak{q}'} \end{array} \left(\begin{array}{cc} \kappa^2(2\delta_{\mathfrak{p}'}^{\mathfrak{p}}\delta_{\mathfrak{q}'}^{\mathfrak{q}} + g^{\mathfrak{p}\mathfrak{q}}g_{\mathfrak{p}'\mathfrak{q}'} - 2\delta_{\mathfrak{q}'}^{\mathfrak{p}}\delta_{\mathfrak{p}'}^{\mathfrak{q}}) & \\ \kappa^2\delta_{\mathcal{J}'}^{\mathcal{I}'}\delta_{\hat{\mathcal{J}}'}^{\hat{\mathcal{I}}'}g^{\mathfrak{p}\mathfrak{q}} & \kappa^2(2\delta_{\hat{\mathcal{I}}}^{\mathcal{I}'}\delta_{\mathcal{J}'}^{\mathcal{J}} - \delta_{\hat{\mathcal{I}}}^{\mathcal{J}}\delta_{\mathcal{J}'}^{\mathcal{I}'})\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}}\delta_{\hat{\mathcal{J}}'}^{\hat{\mathcal{I}}'} + 2\delta_{\hat{\mathcal{I}}}^{\mathcal{J}}\delta_{\mathcal{J}'}^{\mathcal{I}'}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{I}}'}\delta_{\hat{\mathcal{J}}'}^{\hat{\mathcal{J}}} \end{array} \right) \\ & \oplus \begin{array}{cc} \phi^{\mathfrak{p}}Q_{I\hat{I}} & Q_{I\hat{I}}\check{\phi}^{\mathfrak{p}} \\ \phi_{\mathfrak{p}'}\bar{Q}^{\hat{\mathcal{I}}'\mathcal{I}'} & 2\delta_{\hat{\mathcal{I}}}^{\mathcal{I}'}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{I}}'}\delta_{\mathfrak{p}'}^{\mathfrak{p}} & -2\kappa\delta_{\hat{\mathcal{I}}}^{\mathcal{I}'}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{I}}'}\delta_{\mathfrak{p}'}^{\mathfrak{p}} \\ \bar{Q}^{\hat{\mathcal{I}}'\mathcal{I}'}\check{\phi}_{\mathfrak{p}'} & -2\kappa\delta_{\hat{\mathcal{I}}}^{\mathcal{I}'}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{I}}'}\delta_{\mathfrak{p}'}^{\mathfrak{p}} & 2\kappa^2\delta_{\hat{\mathcal{I}}}^{\mathcal{I}'}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{I}}'}\delta_{\mathfrak{p}'}^{\mathfrak{p}} \end{array} \left(\begin{array}{cc} \check{\phi}^{\mathfrak{p}}\bar{Q}^{\hat{\mathcal{I}}\mathcal{J}} & \bar{Q}^{\hat{\mathcal{I}}\mathcal{J}}\phi^{\mathfrak{p}} \\ \check{\phi}_{\mathfrak{p}'}Q_{\mathcal{J}'\hat{\mathcal{J}}'} & 2\kappa^2\delta_{\mathcal{J}'}^{\mathcal{J}}\delta_{\hat{\mathcal{J}}'}^{\hat{\mathcal{J}}}\delta_{\mathfrak{p}'}^{\mathfrak{p}} & -2\kappa\delta_{\mathcal{J}'}^{\mathcal{J}}\delta_{\hat{\mathcal{J}}'}^{\hat{\mathcal{J}}}\delta_{\mathfrak{p}'}^{\mathfrak{p}} \\ Q_{\mathcal{J}'\hat{\mathcal{J}}'}\phi_{\mathfrak{p}'} & -2\kappa\delta_{\mathcal{J}'}^{\mathcal{J}}\delta_{\hat{\mathcal{J}}'}^{\hat{\mathcal{J}}}\delta_{\mathfrak{p}'}^{\mathfrak{p}} & 2\delta_{\mathcal{J}'}^{\mathcal{J}}\delta_{\hat{\mathcal{J}}'}^{\hat{\mathcal{J}}}\delta_{\mathfrak{p}'}^{\mathfrak{p}} \end{array} \right) \end{aligned} \quad (3.16)$$

In concise form,¹⁰

$$H_{k,k+1} = \begin{matrix} & \phi\phi & Q\bar{Q} & \check{\phi}\check{\phi} & \bar{Q}Q & \phi Q & Q\check{\phi} & \check{\phi}\bar{Q} & \bar{Q}\phi \end{matrix} \begin{pmatrix} \phi\phi & (2 + \mathbb{K} - 2\mathbb{P}) & \mathbb{K} & 0 & 0 & 0 & 0 & 0 & 0 \\ Q\bar{Q} & \mathbb{K} & (2 - \mathbb{K})\hat{\mathbb{K}} + 2\kappa^2\mathbb{K} & 0 & 0 & 0 & 0 & 0 & 0 \\ \check{\phi}\check{\phi} & 0 & 0 & \kappa^2(2 + \mathbb{K} - 2\mathbb{P}) & \kappa^2\mathbb{K} & 0 & 0 & 0 & 0 \\ \bar{Q}Q & 0 & 0 & \kappa^2\mathbb{K} & \kappa^2(2 - \mathbb{K})\hat{\mathbb{K}} + 2\mathbb{K} & 0 & 0 & 0 & 0 \\ \phi Q & 0 & 0 & 0 & 0 & 2 & -2\kappa & 0 & 0 \\ Q\check{\phi} & 0 & 0 & 0 & 0 & -2\kappa & 2\kappa^2 & 0 & 0 \\ \check{\phi}\bar{Q} & 0 & 0 & 0 & 0 & 0 & 0 & 2\kappa^2 & -2\kappa \\ \bar{Q}\phi & 0 & 0 & 0 & 0 & 0 & 0 & -2\kappa & 2 \end{pmatrix}$$

where

$$\kappa \equiv \frac{\check{g}}{g}, \quad g^2 \equiv \frac{g_{YM}^2 N}{8\pi^2}, \quad \check{g}^2 \equiv \frac{\check{g}_{YM}^2 N}{8\pi^2}. \quad (3.17)$$

It is easy to check that in the limit $\kappa \rightarrow 0$ this Hamiltonian reduces to that of the SCQCD spin chain, as it should.¹¹

The Hamiltonian can also be compactly written in terms of the $\mathbb{Z}_2$ -projected $SU(2N_c)$ adjoint fields Z and $\mathcal{X}$,

$$Z = \begin{pmatrix} \phi & 0 \\ 0 & \check{\phi} \end{pmatrix}, \quad \mathcal{X}_{\mathcal{I}\hat{\mathcal{I}}} = \begin{pmatrix} 0 & Q_{\mathcal{I}\hat{\mathcal{I}}} \\ -\epsilon_{\mathcal{I}\mathcal{J}}\epsilon_{\hat{\mathcal{I}}\hat{\mathcal{J}}} \bar{Q}^{\hat{\mathcal{J}}\mathcal{J}} & 0 \end{pmatrix}. \quad (3.18)$$

In this notation,

$$g^2 H = \begin{pmatrix} ZZ & \mathcal{X}\mathcal{X} & Z\mathcal{X} & \mathcal{X}Z \\ (g_+ + \gamma g_-)^2(2 + \mathbb{K} - 2\mathbb{P}) & (g_+ + \gamma g_-)^2\mathbb{K}\hat{\mathbb{K}} & 0 & 0 \\ (g_+ + \gamma g_-)^2\mathbb{K}\hat{\mathbb{K}} & (g_+ + \gamma g_-)^2(2\hat{\mathbb{K}} - \mathbb{K}\hat{\mathbb{K}}) + 2(g_+ - \gamma g_-)^2\mathbb{K} & 0 & 0 \\ 0 & 0 & 2(g_+ + \gamma g_-)^2 & -2(g_+^2 - g_-^2) \\ 0 & 0 & -2(g_+^2 - g_-^2) & 2(g_+ - \gamma g_-)^2 \end{pmatrix}, \quad (3.19)$$

where γ is the twist operator (2.11), and we have defined $g_{\pm} \equiv (g \pm \check{g})/2$. The Hamiltonian

¹⁰The meaning of the different operators can be read off by comparing with the explicit form above. Note in particular that to avoid cluttering we have dropped the identity symbol $\mathbb{I}$. Also in the subspaces $Q\bar{Q}$, $\bar{Q}Q$ we use the notation $\mathbb{K}$ for the trace operator acting on $SU(2)_R$ indices and $\hat{\mathbb{K}}$ that acts on the $SU(2)_L$ indices.

¹¹In the comparison, it is important to take into account the factors that arise by normalizing to one the tree-level two-point function. Recall that in SCQCD $\bar{Q}_i Q^i$ is contracted summing over the $N_f = 2N_c$ flavors, while in the interpolating SCFT $\bar{Q}_{\bar{a}} Q^{\bar{a}}$ is contracted summing over the N_c colors (leaving open the $SU(2)_L$ indices).

is invariant under the parity operation¹²

$$Z_b^a \rightarrow -Z_a^b \quad \mathcal{X}_b^a \rightarrow -\mathcal{X}_a^b. \quad (3.20)$$

where here $a, b = 1, \dots, 2N_c$. As it is an exact invariance of the Lagrangian, this parity symmetry is expected to persist to all loops.

3.4 Magnons in the interpolating spin chain

The spin chain of the interpolating SCFT admits two degenerate chiral vacua with $\Delta + r = 0$, namely $\text{Tr } \phi^\ell$ and $\text{Tr } \check{\phi}^\ell$. The elementary impurities are Q and $\bar{Q}$, which have $\Delta + r = 1$. In the infinite chain it makes sense to consider states with a single impurity. A single Q impurity separates the ϕ vacuum to its left from the $\check{\phi}$ vacuum on its right; viceversa for a $\bar{Q}$ impurity.

The action of the Hamiltonian on a single Q impurity in position space is

$$g^2 H Q_{\mathcal{I}\hat{\mathcal{I}}}(x) = 2(g^2 + \check{g}^2) Q_{\mathcal{I}\hat{\mathcal{I}}}(x) - 2g\check{g}[Q_{\mathcal{I}\hat{\mathcal{I}}}(x-1) + Q_{\mathcal{I}\hat{\mathcal{I}}}(x+1)] \quad (3.21)$$

Fourier transforming as $Q(p) = \sum_x e^{ipx} Q(x)$ we have

$$\begin{aligned} g^2 H Q_{\mathcal{I}\hat{\mathcal{I}}}(p) &= 2(g^2 + \check{g}^2 - 2g\check{g} \cos p) Q_{\mathcal{I}\hat{\mathcal{I}}}(p) \\ &= [2(g - \check{g})^2 + 4g\check{g}(1 - \cos p)] Q_{\mathcal{I}\hat{\mathcal{I}}}(p) \\ &= [2(g - \check{g})^2 + 8g\check{g} \sin^2(\frac{p}{2})] Q_{\mathcal{I}\hat{\mathcal{I}}}(p) \end{aligned} \quad (3.22)$$

Hence the dispersion relation for $Q_{\mathcal{I}\hat{\mathcal{I}}}(p)$ is,

$$E(p; \kappa) = 2(1 - \kappa)^2 + 8\kappa \left(\sin^2 \frac{p}{2} \right). \quad (3.23)$$

The magnon is gapless at the orbifold point $\kappa = 1$, and it develops a gap as we move towards SCQCD. Precisely at the SCQCD point, the single impurity state ceases to be meaningful and its dispersion relation trivializes. An identical analysis holds for the $\bar{Q}$ impurity, leading to the same dispersion relation.

4. Protected Spectrum

In this section we put to use the one-loop Hamiltonian to study the protected spectrum of $\mathcal{N} = 2$ SCQCD and of the interpolating SCFT. The results presented here were quoted without proof and used in our previous paper [1]. The remainder of the present paper is independent of this section, and readers mainly interested in dynamics and integrability of the spin chain may proceed directly to section 5.

We are going to determine all the generalized single-trace operators in the scalar sector of SCQCD having vanishing one-loop anomalous dimension. We find the complete list of such

¹²We are indebted to Pedro Liendo for this observation, which corrects the contrary claim made in v1 of the arXiv submission of this paper.

operators to be:¹³

$$\text{Tr } \phi^{k+2}, \quad \text{Tr}[T\phi^k], \quad \text{Tr } \mathcal{M}_3. \quad (4.1)$$

Here, $T \equiv \phi\bar{\phi} - \mathcal{M}_1$ and $k \geq 0$. We are first led to (4.1) by an educated guess. In section 4.1 we list all operators in the scalar sector that obey any of the shortening or semi-shortening conditions of the $\mathcal{N} = 2$ superconformal algebra, which have been completely classified [25, 26, 27, 28, 29]. Using the spin-chain Hamiltonian, we compute the one-loop anomalous dimension of these candidate protected states, and find that only (4.1) have zero anomalous dimension. Even though here we only perform a one-loop analysis, the operators (4.1) can be seen to be protected at full quantum level using the superconformal index [1].

In section 4.2, we list the protected operators of the orbifold theory (they can be exhaustively enumerated by a variety of methods [1]) and follow their evolution along the exactly marginal line κ .

4.1 Protected spectrum in $\mathcal{N} = 2$ SCQCD

A generic long multiplet $\mathcal{A}_{R,r(j,\bar{j})}^\Delta$ of the $\mathcal{N} = 2$ superconformal algebra is generated by the action of the 8 Poincaré supercharges $\mathcal{Q}$ and $\bar{\mathcal{Q}}$ on a superconformal primary, which by definition is annihilated by all conformal supercharges $\mathcal{S}$. If some combination of the $\mathcal{Q}$'s also annihilates the primary, the corresponding multiplet is shorter and the conformal dimensions of all its members are protected against quantum corrections. We follow the conventions of [28] for the possible shortening conditions for the $\mathcal{N} = 2$ superconformal algebra, see table 3.

In table 4 we list all the generalized single-trace operators of $\mathcal{N} = 2$ SCQCD made out of scalar fields, which obey any of the possible shortening conditions. Using the spin-chain Hamiltonian of section 3.1, we find that the only operators with zero anomalous dimension are the one listed in (4.1)¹⁴. The operators $\text{Tr } \phi^\ell$ correspond to the vacuum of the spin chain, while the operators $\text{Tr } T \phi^\ell$ correspond to the zero-momentum limit of the gapless excitation $T(p)$, eq. (3.12). There is one more protected operator, which is “exceptional” in not belonging to an infinite sequence: $\text{Tr } \mathcal{M}_3$. Its anomalous dimension is zero for gauge group $SU(N_c)$ but not for gauge group $U(N_c)$: the double-trace terms in the Lagrangian that arise from the removal of the $U(1)$ are crucial for the protection of this operator (see footnote at page 12).

4.2 Protected spectrum in the orbifold theory

As we have reviewed in section 2.2, $\mathcal{N} = 2$ SCQCD can be obtained as the $\check{g}_{YM} \rightarrow 0$ limit of a family of $\mathcal{N} = 2$ superconformal field theories, which reduces for $g_{YM} = \check{g}_{YM}$ to the $\mathcal{N} = 2$ $\mathbb{Z}_2$ orbifold of $\mathcal{N} = 4$ SYM. In this section we find the protected spectrum of single-trace operators of the interpolating family. We start at the orbifold point, where the protected

¹³As explained in [1], $\mathcal{N} = 2$ SCQCD has a second class of protected operators, which are outside the scalar sector.

¹⁴Together of course with their conjugates. Note that since in our conventions ϕ has $r = -1$, the multiplet $\bar{\mathcal{E}}_{-\ell(0,0)}$, $\ell > 0$, is represented by $\text{Tr } \phi^\ell$. The conjugate multiplet $\mathcal{E}_{\ell(0,0)}$ is represented by $\text{Tr } \bar{\phi}^\ell$ and is of course also protected.

Shortening Conditions				Multiplet
$\mathcal{B}_1$	$\mathcal{Q}_\alpha^1 R, r\rangle^{h.w.} = 0$	$j = 0$	$\Delta = 2R + r$	$\mathcal{B}_{R,r(0,\bar{j})}$
$\bar{\mathcal{B}}_2$	$\bar{\mathcal{Q}}_{2\dot{\alpha}} R, r\rangle^{h.w.} = 0$	$\bar{j} = 0$	$\Delta = 2R - r$	$\bar{\mathcal{B}}_{R,r(j,0)}$
$\mathcal{E}$	$\mathcal{B}_1 \cap \mathcal{B}_2$	$R = 0$	$\Delta = r$	$\mathcal{E}_{r(0,\bar{j})}$
$\bar{\mathcal{E}}$	$\bar{\mathcal{B}}_1 \cap \bar{\mathcal{B}}_2$	$R = 0$	$\Delta = -r$	$\bar{\mathcal{E}}_{r(j,0)}$
$\hat{\mathcal{B}}$	$\mathcal{B}_1 \cap \bar{\mathcal{B}}_2$	$r = 0, j, \bar{j} = 0$	$\Delta = 2R$	$\hat{\mathcal{B}}_R$
$\mathcal{C}_1$	$\epsilon^{\alpha\beta} \mathcal{Q}_\beta^1 R, r\rangle_\alpha^{h.w.} = 0$ $(\mathcal{Q}^1)^2 R, r\rangle^{h.w.} = 0$ for $j = 0$		$\Delta = 2 + 2j + 2R + r$ $\Delta = 2 + 2R + r$	$\mathcal{C}_{R,r(j,\bar{j})}$ $\mathcal{C}_{R,r(0,\bar{j})}$
$\bar{\mathcal{C}}_2$	$\epsilon^{\dot{\alpha}\dot{\beta}} \bar{\mathcal{Q}}_{2\dot{\beta}} R, r\rangle_{\dot{\alpha}}^{h.w.} = 0$ $(\bar{\mathcal{Q}}_2)^2 R, r\rangle^{h.w.} = 0$ for $\bar{j} = 0$		$\Delta = 2 + 2\bar{j} + 2R - r$ $\Delta = 2 + 2R - r$	$\bar{\mathcal{C}}_{R,r(j,\bar{j})}$ $\bar{\mathcal{C}}_{R,r(j,0)}$
$\mathcal{F}$	$\mathcal{C}_1 \cap \mathcal{C}_2$	$R = 0$	$\Delta = 2 + 2j + r$	$\mathcal{C}_{0,r(j,\bar{j})}$
$\bar{\mathcal{F}}$	$\bar{\mathcal{C}}_1 \cap \bar{\mathcal{C}}_2$	$R = 0$	$\Delta = 2 + 2\bar{j} - r$	$\bar{\mathcal{C}}_{0,r(j,\bar{j})}$
$\hat{\mathcal{C}}$	$\mathcal{C}_1 \cap \bar{\mathcal{C}}_2$	$r = \bar{j} - j$	$\Delta = 2 + 2R + j + \bar{j}$	$\hat{\mathcal{C}}_{R(j,\bar{j})}$
$\hat{\mathcal{F}}$	$\mathcal{C}_1 \cap \mathcal{C}_2 \cap \bar{\mathcal{C}}_1 \cap \bar{\mathcal{C}}_2$	$R = 0, r = \bar{j} - j$	$\Delta = 2 + j + \bar{j}$	$\hat{\mathcal{C}}_{0(j,\bar{j})}$
$\mathcal{D}$	$\mathcal{B}_1 \cap \bar{\mathcal{C}}_2$	$r = \bar{j} + 1$	$\Delta = 1 + 2R + \bar{j}$	$\mathcal{D}_{R(0,\bar{j})}$
$\bar{\mathcal{D}}$	$\bar{\mathcal{B}}_2 \cap \mathcal{C}_1$	$-r = j + 1$	$\Delta = 1 + 2R + j$	$\bar{\mathcal{D}}_{R(j,0)}$
$\mathcal{G}$	$\mathcal{E} \cap \bar{\mathcal{C}}_2$	$r = \bar{j} + 1, R = 0$	$\Delta = r = 1 + \bar{j}$	$\mathcal{D}_{0(0,\bar{j})}$
$\bar{\mathcal{G}}$	$\bar{\mathcal{E}} \cap \mathcal{C}_1$	$-r = j + 1, R = 0$	$\Delta = -r = 1 + j$	$\bar{\mathcal{D}}_{0(j,0)}$

Table 3: Shortening conditions and short multiplets for the $\mathcal{N} = 2$ superconformal algebra.

Scalar Multiplets	SCQCD operators	Protected
$\bar{\mathcal{B}}_{R,-\ell(0,0)}$	$\text{Tr}[\phi^\ell \mathcal{M}_3^R]$	
$\mathcal{E}_{-\ell(0,0)}$	$\text{Tr}[\phi^\ell]$	✓
$\bar{\mathcal{B}}_R$	$\text{Tr}[\mathcal{M}_3^R]$	✓ for $R = 1$
$\bar{\mathcal{C}}_{R,-\ell(0,0)}$	$\text{Tr}[T \mathcal{M}_3^R \phi^\ell]$	
$\mathcal{C}_{0,-\ell(0,0)}$	$\text{Tr}[T \phi^\ell]$	✓
$\hat{\mathcal{C}}_{R(0,0)}$	$\text{Tr}[T \mathcal{M}_3^R]$	
$\hat{\mathcal{C}}_{0(0,0)}$	$\text{Tr}[T]$	✓
$\mathcal{D}_{R(0,0)}$	$\text{Tr}[\mathcal{M}_3^R \phi]$	

Table 4: $\mathcal{N} = 2$ SCQCD protected operators at one loop

states are easy to determine, and follow their fate along the exactly marginal line towards $\mathcal{N} = 2$ SCQCD.

At the orbifold point, operators fall into two classes: untwisted and twisted. In the untwisted sector, the protected states are well-known, since they are inherited from $\mathcal{N} = 4$ SYM. The protected operators in the twisted sector are chiral with respect to $\mathcal{N} = 1$ subalgebra and could be obtained by analyzing the chiral ring [30].¹⁵ Both the classes of operators can be rigorously checked to be protected by computing the superconformal index.¹⁶

¹⁵We confirm the spectrum in [1] up to one operator that was missed in the analysis of [30].

¹⁶The calculation for the orbifold was carried out already in [31], and confirmed in [1] up to a minor

Using the index one can also argue that the protected multiplets found at the orbifold point *cannot* recombine into long multiplets as we vary $\check{g}$ [1], so in particular taking $\check{g} \rightarrow 0$ they must evolve into the protected multiplets of the theory

$$\{\mathcal{N} = 2 \text{ SCQCD} \oplus \text{decoupled } SU(N_{\tilde{c}}) \text{ vector multiplet}\}. \quad (4.2)$$

In section 4.3 we follow this evolution in detail. We find that the $SU(2)_L$ -singlet protected states of the interpolating theory evolve into the list (4.1) of protected states of SCQCD, plus some extra states made purely from the decoupled vector multiplet. On the other hand, the interpolating theory has also many single-trace protected states with non-trivial $SU(2)_L$ spin, which are of course absent from the list (4.1): we see that in the limit $\check{g} \rightarrow 0$, a state with $SU(2)_L$ spin L can be interpreted as a “multiparticle state”, obtained by linking together L short “open” spin chains of SCQCD and decoupled fields $\check{\phi}$. By this route we confirm that (4.1) is the correct and complete list of protected single-traces in the scalar sector for $\mathcal{N} = 2$ SCQCD. The results are also suggestive of a dual string theory interpretation: as $\check{g} \rightarrow 0$, single closed string states carrying $SU(2)_L$ quantum numbers disintegrate into multiple open strings. The above argument, however, doesn’t imply that all the protected operators of SCQCD are obtained as degenerations of protected operators of the interpolating theory. Indeed, they aren’t. In [1], we discuss an alternative mechanism that brings about more protected SCQCD operators from the decomposition of long multiplets of the interpolating theory as $\check{g} \rightarrow 0$.

In summary, the degeneracy of protected states is independent of the exactly marginal deformation that changes $\check{g}_{YM}$ and is thus the same for the orbifold theory and for the theory (4.2). At $\check{g}_{YM} = 0$ there is a symmetry enhancement, $SU(2)_L \times SU(N_{\tilde{c}}) \rightarrow U(N_f = 2N_c)$, and we can consistently truncate the spectrum of generalized single trace operators to singlets of the flavor group $U(N_f)$ – which in particular do not contain any of the decoupled states $\check{\phi}$. This is the flavor singlet spectrum of $\mathcal{N} = 2$ SCQCD that we have analyzed in the previous section.

4.3 Away from the orbifold point: matching with $\mathcal{N} = 2$ SCQCD

In the limit $\check{g} \rightarrow 0$, we must be able to match the protected states of the interpolating family with protected states of $\{\mathcal{N} = 2 \text{ SCQCD} \oplus \text{decoupled vector multiplet}\}$. For the purpose of this discussion, the protected states naturally splits into two sets: $SU(2)_L$ singlets and $SU(2)_L$ non-singlets. It is clear that all the (generalized) single-trace operators of $\mathcal{N} = 2$ SCQCD must arise from the $SU(2)_L$ singlets.

emendation.

Multiplet	Orbifold operator ($R, \ell \geq 0, n \geq 2$)
$\hat{\mathcal{B}}_{R+1}$	$\text{Tr}[(Q^{++}\bar{Q}^{++})^{R+1}]$
$\bar{\mathcal{E}}_{-(\ell+2)(0,0)}$	$\text{Tr}[\phi^{\ell+2} + \check{\phi}^{\ell+2}]$
$\hat{\mathcal{C}}_{R(0,0)}$	$\text{Tr}[\sum \mathcal{T}(Q^{++}\bar{Q}^{++})^R]$
$\bar{\mathcal{D}}_{R+1(0,0)}$	$\text{Tr}[\sum (Q^{++}\bar{Q}^{++})^{R+1}(\phi^+\check{\phi})]$
$\bar{\mathcal{B}}_{R+1, -(\ell+2)(0,0)}$	$\text{Tr}[\sum_i (Q^{++}\bar{Q}^{++})^{R+1} \phi^i \check{\phi}^{\ell+2-i}]$
$\bar{\mathcal{C}}_{R, -(\ell+1)(0,0)}$	$\text{Tr}[\sum_i \mathcal{T}(Q^{++}\bar{Q}^{++})^R \phi^i \check{\phi}^{\ell+1-i}]$
$\mathcal{A}_{R, -\ell(0,0)}^{\Delta=2R+\ell+2n}$	$\text{Tr}[\sum_i \mathcal{T}^n(Q^{++}\bar{Q}^{++})^R \phi^i \check{\phi}^{\ell-i}]$

Table 5: Superconformal primary operators in the untwisted sector of the orbifold theory that descend from the $\frac{1}{2}$ BPS primary of $\mathcal{N} = 4$. The symbol $\sum$ indicates summation over all “symmetric traceless” permutations of the component fields allowed by the index structure.

Multiplet	Orbifold operator ($\ell \geq 0$)
$\hat{\mathcal{B}}_1$	$\text{Tr}[(Q^{++}\bar{Q}^{++} - Q^{+-}\bar{Q}^{+-})] = \text{Tr } \mathcal{M}_3$
$\bar{\mathcal{E}}_{-(\ell+2)(0,0)}$	$\text{Tr}[\phi^{\ell+2} - \check{\phi}^{\ell+2}]$

Table 6: Superconformal primary operators in the twisted section of the orbifold theory.

4.3.1 $\text{SU}(2)_{\text{L}}$ singlets

They are:

- (i) One $\hat{\mathcal{B}}_1$ multiplet, corresponding to the primary $\text{Tr}[Q_{\hat{\mathcal{I}}\{\mathcal{I}}}\bar{Q}_{\hat{\mathcal{J}}\}}^{\hat{\mathcal{I}}}] = \text{Tr } \mathcal{M}_3$. Since this is the only operator with these quantum numbers, it cannot mix with anything and its form is independent of $\check{g}$.

- (ii) Two $\bar{\mathcal{E}}_{-\ell(0,0)}$ multiplets for each $\ell \geq 2$, corresponding to the primaries $\text{Tr}[\phi^\ell \pm \check{\phi}^\ell]$.

For each ℓ , there is a two-dimensional space of protected operators, and we may choose whichever basis is more convenient. For $g = \check{g}$, the natural basis vectors are the untwisted and twisted combinations (respectively even and odd under $\phi \leftrightarrow \check{\phi}$), while for $\check{g} = 0$ the natural basis vectors are $\text{Tr } \phi^\ell$ (which is an operator of $\mathcal{N} = 2$ SCQCD) and $\text{Tr } \check{\phi}^\ell$ (which belongs to the decoupled sector).

- (iii) One $\hat{\mathcal{C}}_{0(0,0)}$ multiplet (the stress-tensor multiplet), corresponding to the primary $\text{Tr } \mathcal{T} = \text{Tr } [T + \check{\phi}\check{\phi}]$. We have checked that this combination is an eigenstate with zero eigenvalue for all $\check{g}$.

For $\check{g} = 0$, we may trivially subtract out the decoupled piece $\text{Tr } \check{\phi}\check{\phi}$ and recover $\text{Tr } T$, the stress-tensor multiplet of $\mathcal{N} = 2$ SCQCD.

- (iv) One $\bar{\mathcal{C}}_{0, -\ell(0,0)}$ multiplet for each $\ell \geq 1$. In the limit $\check{g} \rightarrow 0$, we expect this multiplet to evolve to the $\text{Tr } T \phi^\ell$ multiplet of $\mathcal{N} = 2$ SCQCD. Let us check this in detail.

The primary of $\bar{\mathcal{C}}_{0,-\ell(0,0)}$ has $R = 0$, $r = -\ell$ and $\Delta = \ell + 2$. The space of operators which classically have these quantum numbers is spanned by

$$|a\rangle = \text{Tr}[\check{\phi}^{\ell+1}\bar{\phi}], \quad |b_i\rangle \equiv \frac{1}{2}\text{Tr}[\phi^i Q_{\mathcal{I}\hat{\mathcal{I}}} \check{\phi}^{\ell-i} \bar{Q}^{\hat{\mathcal{I}}\mathcal{I}}] \quad \text{for } 0 \leq i \leq \ell \quad \text{and} \quad |c_\ell\rangle \equiv \text{Tr}[\phi^{\ell+1}\bar{\phi}] \quad (4.3)$$

Diagonalizing the Hamiltonian in Fourier space, we find the protected operator to be

$$|\bar{\mathcal{C}}_{0,-\ell(0,0)}\rangle_\kappa = \kappa^\ell |a\rangle - \sum_{i=0}^{\ell} \kappa^{\ell-i} |b_i\rangle + |c_\ell\rangle \quad (4.4)$$

where $\kappa \equiv \check{g}/g$. In the limit $\kappa \rightarrow 0$,

$$|\bar{\mathcal{C}}_{0,-\ell(0,0)}\rangle_{\kappa \rightarrow 0} = \text{Tr}[(\phi\bar{\phi} - \frac{1}{2}Q_{\mathcal{I}\hat{\mathcal{I}}} \bar{Q}^{\hat{\mathcal{I}}\mathcal{I}})\phi^\ell] = \text{Tr}[T\phi^\ell], \quad (4.5)$$

as claimed.

All in all, we see that this list reproduces the list (4.1) of one-loop protected scalar operators of $\mathcal{N} = 2$ SCQCD, *plus* the extra states $\text{Tr}\check{\phi}^\ell$ which decouple for $\check{g} = 0$. This concludes the argument that the operators (4.1) are protected at the full quantum level, and that they are the *complete* set of protected generalized single-trace primaries of $\mathcal{N} = 2$ SCQCD.

4.3.2 $SU(2)_L$ non-singlets

The basic protected primary of $\mathcal{N} = 2$ SCQCD which is charged under $SU(2)_L$ is the $SU(2)_L$ triplet contained in the mesonic operator $\mathcal{O}_{\mathbf{3_R}j}^i = (\bar{Q}_a^i Q_j^a)_{\mathbf{3_R}}$. Indeed writing the $U(N_f = 2N_c)$ flavor indices i as $i = (\check{a}, \hat{\mathcal{I}})$, with $\check{a} = 1, \dots, N_f/2 = N_c$ “half” flavor indices and $\mathcal{I} = \pm$ $SU(2)_L$ indices, we can decompose

$$\mathcal{O}_{\mathbf{3_R}j}^i \rightarrow \mathcal{O}_{\mathbf{3_R} \mathbf{3_L} \check{b}}^{\check{a}}, \quad \mathcal{O}_{\mathbf{3_R} \mathbf{1_L} \check{b}}^{\check{a}}. \quad (4.6)$$

In particular we may consider the highest weight combination for both $SU(2)_L$ and $SU(2)_R$,

$$(\bar{Q}^{+\hat{+}} Q^{+\hat{+}})_{\check{b}}^{\check{a}}. \quad (4.7)$$

States with higher $SU(2)_L$ spin can be built by taking products of $\mathcal{O}_{\mathbf{3_R} \mathbf{3_L}}$ with $SU(2)_L$ and $SU(2)_R$ indices separately symmetrized – and this is the only way to obtain protected states of $\mathcal{N} = 2$ SCQCD charged under $SU(2)_L$ which have finite conformal dimension in the Veneziano limit. It is then a priori clear that a protected primary of the interpolating theory with $SU(2)_L$ spin L must evolve as $\check{g} \rightarrow 0$ into a product of L copies of $(\bar{Q}^{+\hat{+}} Q^{+\hat{+}})$ and of as many additional decoupled scalars $\check{\phi}$ and $\bar{\phi}$ as needed to make up for the correct $U(1)_r$ charge and conformal dimension. It is amusing to follow in more detail this evolution for the various multiplets:

- (i) $\hat{\mathcal{B}}_R$ multiplet.

This is a trivial case, since for each R there is only one operator with the correct quantum numbers, namely

$$|\hat{\mathcal{B}}_R\rangle_\kappa \equiv \text{Tr}[(Q^{+\dagger}\bar{Q}^{+\dagger})^R], \quad (4.8)$$

for all g and $\check{g}$. We have checked that it is indeed an eigenstate of zero eigenvalue for all couplings.

(ii) $\bar{\mathcal{D}}_{R(0,0)}$ multiplet.

The primary of $\bar{\mathcal{D}}_{R(0,0)}$ has $SU(2)_R$ spin equal R , $U(1)_r$ charge $r = -1$ and $\Delta = 2R + 1$. The space of operators which classically have these quantum numbers is two-dimensional, spanned by $\text{Tr}[(Q^{+\dagger}\bar{Q}^{+\dagger})^R\phi]$ and $\text{Tr}[(\bar{Q}^{+\dagger}Q^{+\dagger})^R\check{\phi}]$. The spin chain Hamiltonian in this subspace reads

$$g^2 H_{\bar{\mathcal{D}}} = \begin{pmatrix} 4g^2 & -4g\check{g} \\ -4g\check{g} & 4\check{g}^2 \end{pmatrix} \quad (4.9)$$

The protected operator (eigenvector with zero eigenvalue) is

$$|\bar{\mathcal{D}}_{R(0,0)}\rangle_\kappa \equiv \text{Tr}[\kappa(Q^{+\dagger}\bar{Q}^{+\dagger})^R\phi + (\bar{Q}^{+\dagger}Q^{+\dagger})^R\check{\phi}]. \quad (4.10)$$

For $\kappa = 0$, the protected operator is interpreted as a “multi-particle state” of R open chains of SCQCD and one decoupled scalar $\check{\phi}$. For example for $R = 2$, the operator will be broken into the following gauge-invariant pieces,

$$(\bar{Q}^{+\dagger}Q^{+\dagger})_{\check{b}}^{\check{a}}, \quad (\bar{Q}^{+\dagger}Q^{+\dagger})_{\check{c}}^{\check{b}} \quad \text{and} \quad \check{\phi}_{\check{a}}^{\check{c}}. \quad (4.11)$$

In the limit $\check{g} \rightarrow 0$, the “closed chain” of the interpolating theory effectively breaks into “open chains” of $\{\mathcal{N} = 2 \text{ SCQCD} \oplus \text{decoupled multiplet}\}$, with the rupture points at the contractions of the “half-flavor” indices $\check{a}, \check{b}, \check{c}$.

(iii) $\bar{\mathcal{B}}_{R,r(0,0)}$ multiplet.

Finding the protected multiplet for arbitrary coupling amounts to diagonalizing the spin-chain Hamiltonian of the interpolating theory in the space of operators with quantum numbers R, r and $\Delta = 2R - r$. The dimension of this space increases rapidly with R and r . Let us focus on two simple cases.

case 1: $R = 1, r \equiv -\ell < 0$

In this case, the space is $\ell + 1$ dimensional, spanned by

$$|\psi_i\rangle \equiv \text{Tr}[\phi^i Q^{+\dagger}\check{\phi}^{\ell-i}\bar{Q}^{+\dagger}], \quad i = 0, \dots, \ell. \quad (4.12)$$

The protected operator is found to be

$$|\bar{\mathcal{B}}_{1,-\ell(0,0)}\rangle_\kappa \equiv \sum_{i=0}^{\ell} \kappa^i |\psi_i\rangle \quad (4.13)$$

In our schematic notation of $\sum$, introduced earlier, the same operator would read

$$|\bar{\mathcal{B}}_{1,-\ell(0,0)}\rangle_\kappa = \text{Tr}[\sum_i \kappa^i (Q^{+\hat{+}} \bar{Q}^{+\hat{+}}) \phi^i \check{\phi}^{\ell-i}]. \quad (4.14)$$

Note that at $\kappa = 0$, the $U(1)_r$ charge of the operator is all carried by the decoupled scalars $\check{\phi}$ – there are no ϕ . This is again consistent with the picture of the closed chain disintegrating into open pieces.

case 2: $r = -2$, $R = 2$

The relevant vector space is spanned by the operators

$$\begin{aligned} |0\rangle &= \text{Tr}[\phi \phi Q^{+\hat{+}} \bar{Q}^{+\hat{+}} Q^{+\hat{+}} \bar{Q}^{+\hat{+}}] & |\check{0}\rangle &= \text{Tr}[Q^{+\hat{+}} \check{\phi} \check{\phi} \bar{Q}^{+\hat{+}} Q^{+\hat{+}} \bar{Q}^{+\hat{+}}] \\ |1\rangle &= \text{Tr}[\phi Q^{+\hat{+}} \check{\phi} \bar{Q}^{+\hat{+}} Q^{+\hat{+}} \bar{Q}^{+\hat{+}}] & |\check{1}\rangle &= \text{Tr}[Q^{+\hat{+}} \check{\phi} \bar{Q}^{+\hat{+}} \phi Q^{+\hat{+}} \bar{Q}^{+\hat{+}}] \\ |2\rangle &= \text{Tr}[\phi Q^{+\hat{+}} \bar{Q}^{+\hat{+}} \phi Q^{+\hat{+}} \bar{Q}^{+\hat{+}}] & |\check{2}\rangle &= \text{Tr}[Q^{+\hat{+}} \check{\phi} \bar{Q}^{+\hat{+}} Q^{+\hat{+}} \check{\phi} \bar{Q}^{+\hat{+}}] \end{aligned} \quad (4.15)$$

The Hamiltonian in this subspace is (the basis vectors are read in the sequence $|0\rangle, |\check{0}\rangle, |1\rangle, \dots$)

$$g^2 H_{\bar{\mathcal{B}}_{2,-2(0,0)}} = \begin{pmatrix} 4g^2 & 0 & -2g\check{g} & -2g\check{g} & 0 & 0 \\ 0 & 4\check{g}^2 & -2g\check{g} & -2g\check{g} & 0 & 0 \\ -2g\check{g} & -2g\check{g} & 4g^2 + 4\check{g}^2 & 0 & -2g\check{g} & -2g\check{g} \\ -2g\check{g} & -2g\check{g} & 0 & 4g^2 + 4\check{g}^2 & -2g\check{g} & -2g\check{g} \\ 0 & 0 & -2g\check{g} & -2g\check{g} & 4g^2 & 0 \\ 0 & 0 & -2g\check{g} & -2g\check{g} & 0 & 4\check{g}^2 \end{pmatrix} \quad (4.16)$$

There is an eigenvector with zero eigenvalue for all κ , namely

$$\begin{aligned} |\bar{\mathcal{B}}_{2,-2(0,0)}\rangle_\kappa &\equiv \kappa^2 |0\rangle + |\check{0}\rangle + \kappa |1\rangle + \kappa |\check{1}\rangle + \kappa^2 |2\rangle + |\check{2}\rangle \\ &= \text{Tr}[\sum_i \kappa^i (Q^{+\hat{+}} \bar{Q}^{+\hat{+}})^2 \phi^i \check{\phi}^{2-i}] \end{aligned}$$

As expected, for $\kappa = 0$ the operator contains $\check{\phi}$ and no ϕ .

Extrapolating from these cases, we make an educated guess for the form for general protected operator,

$$|\bar{\mathcal{B}}_{R,-\ell(0,0)}\rangle_\kappa = \text{Tr}[\sum_i \kappa^i (Q^{+\hat{+}} \bar{Q}^{+\hat{+}})^R \phi^i \check{\phi}^{\ell-i}]. \quad (4.17)$$

In the limit $\kappa \rightarrow 0$, this operator breaks into R mesons $(\bar{Q}Q)_{\check{b}}^{\check{a}}$ of $\mathcal{N} = 2$ SCQCD and ℓ decoupled scalars $\check{\phi}_{\check{b}}^{\check{a}}$.

(iv) $\hat{\mathcal{C}}_{R(0,0)}$ and $\bar{\mathcal{C}}_{R,-\ell(0,0)}$ multiplets.

We have not studied these cases in detail since they are technically quite involved. It is clear however that for $\check{g} \rightarrow 0$ the protected primaries must evolve into states of the schematic form

$$\text{Tr} \left[\mathcal{O}_{\mathbf{3_R} \mathbf{3_L}}^R \check{\phi}^{\ell+n} \bar{\phi}^n \right], \quad (4.18)$$

with $\ell = 0$, $n = 1$ for $\hat{\mathcal{C}}_{R(0,0)}$ and $n = 1$ for $\bar{\mathcal{C}}_{R,-\ell(0,0)}$.

5. Two-body scattering

In this section we study the scattering of two magnons in the spin chain for the interpolating SCFT. We take the chain to be infinite. Because of the index structure of the impurities, one of the asymptotic magnons must be a Q and the other a $\bar{Q}$, and their ordering is fixed – we can have a Q impurity always to the left of a $\bar{Q}$ impurity, or viceversa. The scattering is thus pure reflection. For the case of Q to the *left* of $\bar{Q}$, and suppressing momentarily the $SU(2)_L \times SU(2)_R$ quantum numbers, the asymptotic form of the eigenstates of the Hamiltonian is

$$\sum_{x_1 \ll x_2} \left(e^{ip_1 x_1 + ip_2 x_2} + S(p_2, p_1) e^{ip_2 x_1 + ip_1 x_2} \right) | \dots \phi Q(x_1) \check{\phi} \dots \check{\phi} \bar{Q}(x_2) \phi \dots \rangle. \quad (5.1)$$

This is the definition of the two-body S -matrix. In fact, thanks to the nearest-neighbor nature of the spin chain, if the impurities are not adjacent we are already in the “asymptotic” region, so $x_1 \ll x_2$ should be interpreted as $x_1 < x_2 - 1$. Similarly, for the case where Q to the *right* of $\bar{Q}$ the asymptotic form of the two-magnon state is

$$\sum_{x_1 \ll x_2} \left(e^{ip_1 x_1 + ip_2 x_2} + \check{S}(p_2, p_1) e^{ip_2 x_1 + ip_1 x_2} \right) | \dots \check{\phi} \bar{Q}(x_1) \phi \dots \phi Q(x_2) \check{\phi} \dots \rangle, \quad (5.2)$$

which defines $\check{S}$. The two-body S -matrices S and $\check{S}$ are related by exchanging $g \leftrightarrow \check{g}$,

$$S(p_1, p_2; g, \check{g}) = \check{S}(p_1, p_2; \check{g}, g). \quad (5.3)$$

The total energy of a two-magnon state is just the sum of the energy of the individual magnons,

$$E(p_1, p_2; \kappa) = \left(2(1 - \kappa)^2 + 8\kappa(\sin^2 \frac{p_1}{2}) \right) + \left(2(1 - \kappa)^2 + 8\kappa(\sin^2 \frac{p_2}{2}) \right). \quad (5.4)$$

Besides the continuum of states with real momenta p_1 and p_2 , there can be bound and “anti-bound” states for special complex values of the momenta. A bound state occurs when

$$S(p_1, p_2) = \infty, \quad \text{with} \quad p_1 = \frac{P}{2} - iq, \quad p_2 = \frac{P}{2} + iq, \quad q > 0. \quad (5.5)$$

Since $S(p_2, p_1) = 1/S(p_1, p_2) \rightarrow 0$, the asymptotic wave-function is

$$e^{iP \frac{x_1 + x_2}{2} - q(x_2 - x_1)}, \quad (5.6)$$

which is indeed normalizable (since $x_2 > x_1$ in our conventions). A bound state has smaller energy than any state in the two-particle continuum with the same total momentum P . An anti-bound state occurs when

$$S(p_1, p_2) = \infty, \quad \text{with} \quad p_1 = \frac{P}{2} - iq + \pi, \quad p_2 = \frac{P}{2} + iq - \pi, \quad q > 0. \quad (5.7)$$

The asymptotic wave-function is now

$$(-1)^{x_2 - x_1} e^{iP \frac{x_1 + x_2}{2} - q(x_2 - x_1)}. \quad (5.8)$$

The energy of an anti-bound state is strictly bigger than the two-particle continuum. It is easy to see that (5.5) and (5.7) are the only allowed possibilities for complex p_1 and p_2 such that the total momentum and the total energy are real.

The analysis of two-body scattering proceeds independently in four different sectors, corresponding the choice of the triplet or singlet combinations for $SU(2)_L$ and $SU(2)_R$. In each sector, we will compute the S-matrix and look for the (anti)bound states associated to its poles.

5.1 $3_L \otimes 3_R$ Sector

In the $3_L \otimes 3_R$ sector, we write the general two-impurity state with Q to the left of $\bar{Q}$ as

$$|\Psi_{3\otimes 3}\rangle = \sum_{x_1 < x_2} \Psi_{3\otimes 3}(x_1, x_2) |\dots \phi Q(x_1) \check{\phi} \dots \check{\phi} \bar{Q}(x_2) \phi \dots\rangle_{3\otimes 3}. \quad (5.9)$$

There is no mixing with states containing $\bar{\phi}$ and $\check{\bar{\phi}}$ since they have different $SU(2)_L \times SU(2)_R \times U(1)_r$ quantum numbers. Acting with the Hamiltonian, one finds:

- For $x_2 > x_1 + 1$,

$$\begin{aligned} g^2 H \cdot \Psi_{3\otimes 3}(x_1, x_2) &= 4(g^2 + \check{g}^2) \Psi_{3\otimes 3}(x_1, x_2) - 2g\check{g} \Psi_{3\otimes 3}(x_1 + 1, x_2) - 2g\check{g} \Psi_{3\otimes 3}(x_1 - 1, x_2) \\ &\quad - 2g\check{g} \Psi_{3\otimes 3}(x_1, x_2 + 1) - 2g\check{g} \Psi_{3\otimes 3}(x_1, x_2 - 1). \end{aligned} \quad (5.10)$$

- For $x_2 = x_1 + 1$,

$$g^2 H \cdot \Psi_{3\otimes 3}(x_1, x_2) = 4g^2 \Psi_{3\otimes 3}(x_1, x_2) - 2g\check{g} \Psi_{3\otimes 3}(x_1 - 1, x_2) - 2g\check{g} \Psi_{3\otimes 3}(x_1, x_2 + 1). \quad (5.11)$$

The plane wave states $e^{i(p_1 x_1 + p_2 x_2)}$ and $e^{i(p_1 x_2 + p_2 x_1)}$ are separately eigenstates for the “bulk” action of the Hamiltonian (5.10), with eigenvalue (5.4). The action of the Hamiltonian on the state with adjacent impurities, equ.(5.11), provides the boundary condition that fixes the exact eigenstate of asymptotic momenta p_1, p_2 ,

$$\Psi_{3\otimes 3}(x_1, x_2) = e^{i(p_1 x_1 + p_2 x_2)} + S_{3\otimes 3}(p_1, p_2) e^{i(p_1 x_2 + p_2 x_1)}, \quad (5.12)$$

where

$$S_{3\otimes 3}(p_1, p_2) = -\frac{1 + e^{ip_1+ip_2} - 2\kappa e^{ip_1}}{1 + e^{ip_1+ip_2} - 2\kappa e^{ip_2}}, \quad \kappa \equiv \frac{\check{g}}{g}. \quad (5.13)$$

In this sector, the S-matrix coincides with the familiar S-matrix of the XXZ chain, with the identification $\Delta_{XXZ} = \kappa$. The pole of the S-matrix,

$$e^{ip_2} = \frac{1 + e^{i(p_1+p_2)}}{2\kappa}, \quad (5.14)$$

is associated to a bound state. Writing $p_1 = P/2 - iq$, $p_2 = P/2 + iq$, we have

$$e^{-q} = \frac{\cos(\frac{P}{2})}{\kappa}. \quad (5.15)$$

The wave-function is normalizable provided $q > 0$, which implies $2\arccos \kappa < |P| < \pi$. Substituting p_1 and p_2 into the expression for the total energy (5.4), we find that the dispersion relation of the bound state is simply

$$[Q\bar{Q}]_{3_L 3_R}^{bound} : \quad E = 4\sin^2(\frac{P}{2}), \quad 2\arccos \kappa < |P| < \pi. \quad (5.16)$$

This dispersion relation is plotted as the dotted (orange) curve in the left column of figure 2. When the total momentum P is smaller than $2\arccos \kappa$ the bound state dissolves into the two-particle continuum. The bound state exists for the full range of P at the orbifold point $\kappa = 1$, but the allowed range of P shrinks as κ is decreased, and the bound state disappears in the SCQCD limit $\kappa \rightarrow 0$.

The S-matrix in the $3_L \otimes 3_R$ sector with Q to the *right* of $\bar{Q}$ is obtained by switching $g \leftrightarrow \check{g}$,

$$\check{S}_{3\otimes 3}(p_1, p_2; \kappa) = S_{3\otimes 3}(p_1, p_2; 1/\kappa) = -\frac{1 + e^{ip_1+ip_2} - \frac{2}{\kappa}e^{ip_1}}{1 + e^{ip_1+ip_2} - \frac{2}{\kappa}e^{ip_2}}. \quad (5.17)$$

Now the pole of the S-matrix is associated to a bound state with

$$e^{-q} = \kappa \cos(\frac{P}{2}). \quad (5.18)$$

The bound state exists for all P in the whole range of $\kappa \in (0, 1]$. Its dispersion relation is

$$[\bar{Q}Q]_{3_L 3_R}^{bound} : \quad E = 4\kappa^2 \sin^2(\frac{P}{2}), \quad (5.19)$$

plotted as the dotted (orange) curve in the right column of figure 2. The existence of this bound state is consistent with our analysis of the protected spectrum in section 4.

5.2 $1_L \otimes 3_R$ Sector

The general two-body state with Q to the left of $\bar{Q}$ is

$$|\Psi_{1\otimes 3}\rangle = \sum_{x_1 < x_2} \Psi_{1\otimes 3}(x_1, x_2) |\dots \phi Q(x_1) \check{\phi} \dots \check{\phi} \bar{Q}(x_2) \phi \dots\rangle_{1\otimes 3} \quad (5.20)$$

The action of the Hamiltonian for $x_2 = x_1 + 1$ is now

$$g^2 H \cdot \Psi_{1\otimes 3}(x, x+1) = 8g^2 \Psi_{1\otimes 3}(x, x+1) - 2g\check{g} \Psi_{1\otimes 3}(x-1, x+1) - 2g\check{g} \Psi_{1\otimes 3}(x, x+2). \quad (5.21)$$

Writing

$$\Psi_{1\otimes 3}(x_1, x_2) = e^{i(p_1 x_1 + p_2 x_2)} + S_{1\otimes 3}(p_2, p_1) e^{i(p_1 x_2 + p_2 x_1)}, \quad (5.22)$$

we find

$$S_{1\otimes 3}(p_1, p_2; \kappa) = -\frac{1 + e^{ip_1 + ip_2} - 2(\kappa - \frac{1}{\kappa})e^{ip_1}}{1 + e^{ip_1 + ip_2} - 2(\kappa - \frac{1}{\kappa})e^{ip_2}}, \quad (5.23)$$

which is again the S-matrix of the XXZ chain, now with $\Delta = \kappa - \frac{1}{\kappa}$. The S-matrix blows up for

$$e^{ip_2} = \frac{1 + e^{i(p_1 + p_2)}}{2(\kappa - \frac{1}{\kappa})}. \quad (5.24)$$

This pole is associated to an *anti*-bound state. Parametrizing $p_1 = P/2 - iq + \pi$, $p_2 = P/2 - iq - \pi$, the location of the pole is given by

$$e^{-q} = \frac{\cos(\frac{P}{2})}{\frac{1}{\kappa} - \kappa}. \quad (5.25)$$

Normalizability of the wave-function requires $q > 0$, which occurs for a restricted range of P for $\kappa_* < \kappa < 1$, and for the full range of P for $\kappa < \kappa_*$,

$$\begin{aligned} 2 \arccos(\frac{1}{\kappa} - \kappa) < |P| < \pi & \text{ for } \frac{\sqrt{5}-1}{2} < \kappa < 1 \\ 0 < |P| < \pi & \text{ for } 0 < \kappa < \frac{\sqrt{5}-1}{2}. \end{aligned} \quad (5.26)$$

Substituting in $E(p_1, p_2; \kappa)$ we find the dispersion relation for the anti-bound state,

$$[Q\bar{Q}]_{1_L 3_R}^{antibound} : \quad E = \frac{4(2 - \kappa^2)}{1 - \kappa^2} - \frac{4\kappa^2}{1 - \kappa^2} \sin^2 \frac{P}{2}, \quad (5.27)$$

which is plotted as the solid (red) curve in the left column of figure 2. The anti-bound state is absent at the orbifold point $\kappa = 1$. For $\kappa \rightarrow 0$, $q \rightarrow +\infty$, so that the wave-function (5.8) localizes to two neighboring sites and in fact coincides with the dimeric excitation $\mathcal{M}_3 = (Q\bar{Q})_3$ of $\mathcal{N} = 2$ SCQCD; in the limit we smoothly recover the $\mathcal{M}_3$ dispersion relation $E(P) = 8$.

For $\bar{Q}Q$ scattering, we have

$$\check{S}_{1\otimes 3}(p_1, p_2; \kappa) = S_{1\otimes 3}(p_1, p_2; 1/\kappa) = -\frac{1 + e^{ip_1+ip_2} - 2(\frac{1}{\kappa} - \kappa)e^{ip_1}}{1 + e^{ip_1+ip_2} - 2(\frac{1}{\kappa} - \kappa)e^{ip_2}}. \quad (5.28)$$

Now the pole corresponds to a bound state, indeed it occurs for $p_1 = P/2 - iq$, $p_2 = P/2 + iq$ with q and P related as in (5.25). Clearly the allowed range of P is as in (5.26). We find the dispersion relation

$$[Q\bar{Q}]_{1_L 3_R}^{bound} : \quad E = \frac{4\kappa^2}{(1 - \kappa^2)}(1 - 2\kappa^2 + \sin^2 \frac{P}{2}), \quad (5.29)$$

which is plotted as the solid (red) curve in the right column of figure 2.

5.3 $3_L \otimes 1_R$ Sector

The scattering problem in the $3_L \otimes 1_R$ sector is solved by the same technique. We find

$$S_{3\otimes 1}(p_1, p_2) = \check{S}_{3\otimes 1}(p_1, p_2) = -1, \quad (5.30)$$

which coincides with the scattering matrix of free fermions, or with the $\Delta_{XXZ} \rightarrow \infty$ limit of the S-matrix for the XXZ chain. Clearly there are no (anti)bound states.

5.4 $1_L \otimes 1_R$ Sector

The analysis for the $1_L \otimes 1_R$ sector is slightly more involved because a two-impurity state is not closed under the action of Hamiltonian: when Q and $\bar{Q}$ are next to each other they can transform into $\phi\bar{\phi}$. The general state for $Q\bar{Q}$ scattering in the singlet sector is

$$\begin{aligned} |\Psi_{1\otimes 1}\rangle = & \sum_{x_1 < x_2} \Psi_{1\otimes 1}(x_1, x_2) |\dots \phi Q(x_1) \check{\phi} \dots \check{\phi} \bar{Q}(x_2) \phi \dots\rangle_{1\otimes 1} \\ & + \sum_x \Psi_{\bar{\phi}}(x) |\dots \phi \bar{\phi}(x) \phi \dots\rangle. \end{aligned} \quad (5.31)$$

The first term is an eigenstate for “bulk” action of the Hamiltonian ($x_2 > x_1 + 1$) with the usual eigenvalue $E(p_1, p_2; \kappa)$ of equ.(5.4). The action of the Hamiltonian for $x_2 = x_1 + 1$ is

$$\begin{aligned} g^2 H \cdot \Psi_{1\otimes 1}(x, x+1) = & 4(g^2 + \check{g}^2) \Psi_{1\otimes 1}(x, x+1) - 2g\check{g} \Psi_{1\otimes 1}(x-1, x+1) - 2g\check{g} \Psi_{1\otimes 1}(x, x+2) \\ & + 2g^2 \Psi_{\bar{\phi}}(x) + 2g^2 \Psi_{\bar{\phi}}(x+1). \end{aligned} \quad (5.32)$$

Furthermore,

$$\begin{aligned} g^2 H \cdot \Psi_{\bar{\phi}}(x) = & 6g^2 \Psi_{\bar{\phi}}(x) - g^2 \Psi_{\bar{\phi}}(x+1) - g^2 \Psi_{\bar{\phi}}(x-1) \\ & + 2g^2 \Psi_{1\otimes 1}(x, x+1) + 2g^2 \Psi_{1\otimes 1}(x-1, x). \end{aligned} \quad (5.33)$$

We take the ansatz

$$\Psi_{1\otimes 1}(x_1, x_2) = e^{i(p_1 x_1 + p_2 x_2)} + S_{1\otimes 1}(p_2, p_1) e^{i(p_1 x_2 + p_2 x_1)} \quad (5.34)$$

$$\Psi_{\bar{\phi}}(x) = S_{\bar{\phi}}(p_2, p_1) e^{i(p_1 + p_2)x}. \quad (5.35)$$

Note that $S_{1\otimes 1}(p_1, p_2)$ still has the interpretation of the scattering matrix of the magnons Q and $\bar{Q}$, which are the asymptotic excitations, while $\bar{\phi}$ is an “unstable” excitations created during the collision of Q and $\bar{Q}$. We find

$$S_{1\otimes 1}(p_1, p_2) = - \left(\frac{1 + e^{ip_1 + ip_2} - 2(\kappa - \frac{1}{\kappa})e^{ip_1}}{1 + e^{ip_1 + ip_2} - 2(\kappa - \frac{1}{\kappa})e^{ip_2}} \right) \left(\frac{1 + e^{ip_1 + ip_2} - 2\kappa e^{ip_1}}{1 + e^{ip_1 + ip_2} - 2\kappa e^{ip_2}} \right)^{-1} \quad (5.36)$$

$$S_{\bar{\phi}}(p_1, p_2) = \frac{4ie^{i(p_1 + p_2)}(\sin p_1 - \sin p_2)}{(1 + e^{ip_1 + ip_2} - 2\kappa e^{ip_1})(1 + e^{ip_1 + ip_2} - 2(\kappa - \frac{1}{\kappa})e^{ip_2})}. \quad (5.37)$$

$S_{1\otimes 1}$ is the product of two factors, and it admits two poles. The first factor coincides with $S_{1\otimes 3}$, so its pole is associated to an anti-bound state entirely analogous to the anti-bound state in the $1_L \otimes 3_R$ sector. The pole is located at $p_1 = P/2 - iq + \pi$, $p_2 = P/2 + iq - \pi$, with

$$e^{-q} = \frac{\cos(P/2)}{\frac{1}{\kappa} - \kappa}. \quad (5.38)$$

The dispersion relation is again

$$[Q\bar{Q}]_{1_L 1_R}^{antibound} : \quad E = \frac{4(2 - \kappa^2)}{1 - \kappa^2} - \frac{4\kappa^2}{1 - \kappa^2} \sin^2 \frac{P}{2}, \quad (5.39)$$

and the range of P for which the wave-function is normalizable is as in (5.26) – see the solid (red) curve in the left column of figure 2. It is interesting to analyze the explicit form of the wave-function in the $\kappa \rightarrow 0$ limit. The $Q\bar{Q}$ piece has the form

$$\Psi_{1\otimes 1}(x_1, x_2) = (-1)^{x_2 - x_1} e^{iP(\frac{x_1 + x_2}{2})} e^{-q(x_2 - x_1)}, \quad q \rightarrow \infty \quad (5.40)$$

so only the $x_2 = x_1 + 1$ term survives in the limit, and we recover the dimeric impurity $\mathcal{M}_1$ of SCQCD. A short calculation gives

$$\frac{\Psi_{\bar{\phi}}(x)}{\Psi(x, x+1)}|_{\kappa \rightarrow 0} = \frac{2}{(1 + e^{iP})}. \quad (5.41)$$

Comparison with (3.11) shows that that in the $\kappa \rightarrow 0$ limit the antibound state in the $Q\bar{Q}$ singlet sector becomes precisely the dimeric excitation $\tilde{T}$ of SCQCD.

The pole in the second factor of $S_{1\otimes 1}$ corresponds instead to a bound state, with

$$e^q = \frac{\cos(P/2)}{\kappa}. \quad (5.42)$$

The dispersion relation and range of existence are

$$[Q\bar{Q}]_{1_L 1_R}^{bound} : \quad E = 4 \sin^2 \frac{q}{2}, \quad 0 < |P| < 2 \arccos \kappa, \quad (5.43)$$

which are shown as the dashed (green) curve on the left column of figure 2. This bound state is absent at the orbifold point and comes into full existence (for any P) in the SCQCD limit $\kappa \rightarrow 0$. The natural guess is that in this limit it reduces to the gapless $T(p)$ magnon of SCQCD, and it does:

$$\frac{\Psi_{\bar{\phi}}(x)}{\Psi(x, x+1)}|_{\kappa \rightarrow 0} = -\frac{1 + e^{-iP}}{2}, \quad (5.44)$$

in agreement with (3.10).

The S-matrix in the $\bar{Q}Q$ channel is obtained as usual by $\kappa \rightarrow 1/\kappa$,

$$\begin{aligned} \check{S}_{1 \otimes 1}(p_1, p_2; \kappa) &= - \left(\frac{1 + e^{ip_1+ip_2} + 2(\kappa - \frac{1}{\kappa})e^{ip_1}}{1 + e^{ip_1+ip_2} + 2(\kappa - \frac{1}{\kappa})e^{ip_2}} \right) \left(\frac{1 + e^{ip_1+ip_2} - \frac{2}{\kappa}e^{ip_1}}{1 + e^{ip_1+ip_2} - \frac{2}{\kappa}e^{ip_2}} \right)^{-1} \\ \check{S}_{\bar{\phi}}(p_1, p_2; \kappa) &= \frac{4ie^{i(p_1+p_2)}(\sin p_1 - \sin p_2)}{(1 + e^{ip_1+ip_2} - \frac{2}{\kappa}e^{ip_1})(1 + e^{ip_1+ip_2} + 2(\kappa - \frac{1}{\kappa})e^{ip_2})}. \end{aligned}$$

The pole in the first factor of $\check{S}_{1 \otimes 1}$ corresponds to a bound state, with

$$[\bar{Q}Q]_{1_L 1_R}^{bound} : \quad E(P) = \frac{4\kappa^2}{1 - \kappa^2} \left(1 - 2\kappa^2 + \sin^2 \frac{P}{2} \right), \quad (5.45)$$

with the range of existence given by (5.26). Finally, the pole in the second factor of $\check{S}_{1 \otimes 1}$ never corresponds to a normalizable solution.

5.5 Summary

The two-body scattering problem in the spin chain of the interpolating SCFT admits a rich spectrum of bound and anti-bound states. The results are summarized in table 7 and figure 2. The $Q\bar{Q}$ scattering channel (that is, the channel with Q to the left of $\bar{Q}$, and the ϕ vacuum on the outside) is the one relevant to make contact with $\mathcal{N} = 2$ SCQCD, which is obtained in the $\kappa \rightarrow 0$ limit. Remarkably, the magnon excitations of SCQCD are recovered as the smooth limits of the $Q\bar{Q}$ (anti)bound states: as $\kappa \rightarrow 0$ the wavefunctions of the (anti)bound states localize to two neighboring sites and reproduce the “dimeric” magnons $T(p)$, $\tilde{T}(p)$ and $\mathcal{M}_3(p)$ of SCQCD.

5.6 Left/right factorization of the two-body S-matrix

As is well-known, the magnon excitations of the $\mathcal{N} = 4$ SYM spin chain transform in the fundamental representation of $SU(2|2) \times SU(2|2)$, and their two-body S-matrix factorizes into the product of the S-matrices for the “left” and “right” $SU(2|2)$. The $\mathbb{Z}_2$ orbifold preserves this factorization. Remarkably, this left/right factorization persists even away from the orbifold

	Pole of the S-matrix	Range of existence	Dispersion relation $E(P)$
$\mathcal{M}_{33}$	$e^{-q} = \cos(\frac{P}{2})/\kappa$	$2 \arccos \kappa < P < \pi$	$4 \sin^2(\frac{P}{2})$
T	$e^q = \cos(\frac{P}{2})/\kappa$	$0 < P < 2 \arccos \kappa$	$4 \sin^2(\frac{P}{2})$
$\tilde{T}$ and $\mathcal{M}_3$	$e^{-q} = \cos(\frac{P}{2})/(\kappa - \frac{1}{\kappa})$	See equ.(5.26)	$\frac{4\kappa^2}{(1-\kappa^2)}(\frac{2}{\kappa^2} - 1 - \sin^2 \frac{P}{2})$
$\tilde{\mathcal{M}}_{33}$	$e^{-q} = \kappa \cos(\frac{P}{2})$	$0 < P < \pi$	$4\kappa^2 \sin^2(\frac{P}{2})$
$\tilde{T}$	$e^q = \kappa \cos(\frac{P}{2})$	No solution	
$\tilde{\tilde{T}}$ and $\tilde{\mathcal{M}}_3$	$e^{-q} = \cos(\frac{P}{2})/(\frac{1}{\kappa} - \kappa)$	See equ.(5.26)	$\frac{4\kappa^2}{(1-\kappa^2)}(1 - 2\kappa^2 + \sin^2 \frac{P}{2})$

Table 7: Dispersion relations and range of existence of the various (anti)bound states in two-body scattering. The first three entries correspond to the $Q\bar{Q}$ channel and the last three entries to the $\bar{Q}Q$ channel. The color-coding of the third entry is a reminder that these are *anti*-bound states with energy above the two-particle continuum.

point, for the full interpolation SCFT – or at least this is what happens at one-loop in the scalar sector. Our results for the S-matrix in the $Q\bar{Q}$ channel in the four different $SU(2)_L \times SU(2)_R$ sectors are summarized in table 8, where we have defined

$$\mathcal{S}(p_1, p_2, \kappa) \equiv -\frac{1 - 2\kappa e^{ip_1} + e^{i(p_1+p_2)}}{1 - 2\kappa e^{ip_2} + e^{i(p_1+p_2)}}, \quad (5.46)$$

i.e. the standard S-matrix of the XXZ chain, with $\Delta_{X X Z} = \kappa$.

$L \otimes R$	$\mathcal{S}(p_1, p_2, \kappa)$
$1 \otimes 1$	$-\mathcal{S}(p_1, p_2, \kappa - \frac{1}{\kappa})\mathcal{S}^{-1}(p_1, p_2, \kappa)$
$1 \otimes 3$	$\mathcal{S}(p_1, p_2, \kappa - \frac{1}{\kappa})$
$3 \otimes 1$	-1
$3 \otimes 3$	$\mathcal{S}(p_1, p_2, \kappa)$

Table 8: The S-matrix in the $Q\bar{Q}$ scattering channel.

We see that we can write

$$S(p_1, p_2; \kappa) = \frac{S_L(p_1, p_2; \kappa)S_R(p_1, p_2; \kappa)}{S_{3 \otimes 3}(p_1, p_2; \kappa)} \quad (5.47)$$

where S_L and S_R are defined in table 9.

$SU(2)_L$	$S_L(p_1, p_2; \kappa)$	$SU(2)_R$	$S_R(p_1, p_2; \kappa)$
1	$\mathcal{S}(p_1, p_2; \kappa - \frac{1}{\kappa})$	1	-1
3	$\mathcal{S}(p_1, p_2; \kappa)$	3	$\mathcal{S}(p_1, p_2; \kappa)$

Table 9: Definitions of the $SU(2)_L$ and $SU(2)_R$ S-matrices.

In the analysis of the Yang-Baxter equation, it will be useful to write the S-matrices in




Figure 2: Plots of the dispersion relations of the (anti)bound states for different values of κ . The shaded region represents the two-particle continuum.

both the $SU(2)_L$ and $SU(2)_R$ sectors using the identity ($\mathbb{I}$) and trace ($\mathbb{K}$) tensorial structures,

$$S_L(p_1, p_2; \kappa) = A_L(p_1, p_2; \kappa) \mathbb{I} + B_L(p_1, p_2; \kappa) \mathbb{K} \quad (5.48)$$

$$S_R(p_1, p_2; \kappa) = A_R(p_1, p_2; \kappa) \mathbb{I} + B_R(p_1, p_2; \kappa) \mathbb{K}. \quad (5.49)$$

Writing the indices explicitly,

$$(S_R)_{\mathcal{IJ}}^{\mathcal{MN}} = A_R \delta_{\mathcal{I}}^{\mathcal{M}} \delta_{\mathcal{J}}^{\mathcal{N}} + B_R \epsilon_{\mathcal{IJ}} \epsilon^{\mathcal{MN}}, \quad (5.50)$$

Recalling that eigenvalue of $\mathbb{K}$ on the triplet is zero while it is two on the singlet, we see that

$$A = S_{\mathbf{3}} \quad (5.51)$$

$$B = \frac{1}{2}(S_1 - S_{\mathbf{3}}). \quad (5.52)$$

The values of S_1 and $S_{\mathbf{3}}$ in both the $SU(2)_L$ and $SU(2)_R$ sectors can be read off from table 9,

$$A_L(p_1, p_2, \kappa) = \mathcal{S}(p_1, p_2, \kappa) \quad (5.53)$$

$$B_L(p_1, p_2, \kappa) = \frac{1}{2} \left(\mathcal{S}(p_1, p_2, \kappa - \frac{1}{\kappa}) - \mathcal{S}(p_1, p_2, \kappa) \right) \quad (5.54)$$

$$A_R(p_1, p_2, \kappa) = \mathcal{S}(p_1, p_2, \kappa) \quad (5.55)$$

$$B_R(p_1, p_2, \kappa) = -\frac{1}{2}(1 + \mathcal{S}(p_1, p_2, \kappa)). \quad (5.56)$$

In complete analogy, in the $\bar{Q}Q$ channel we have the factorization

$$\check{S}(p_1, p_2; \kappa) = \frac{\check{S}_L(p_1, p_2; \kappa) \check{S}_R(p_1, p_2; \kappa)}{\check{S}_{3 \otimes 3}(p_1, p_2; \kappa)}, \quad (5.57)$$

and we can write

$$\check{S}_L(p_1, p_2; \kappa) = \check{A}_L(p_1, p_2; \kappa) \mathbb{I} + \check{B}_L(p_1, p_2; \kappa) \mathbb{K} \quad (5.58)$$

$$\check{S}_R(p_1, p_2; \kappa) = \check{A}_R(p_1, p_2; \kappa) \mathbb{I} + \check{B}_R(p_1, p_2; \kappa) \mathbb{K}. \quad (5.59)$$

As always, each “checked” quantity is obtained from the corresponding unchecked one by sending $\kappa \rightarrow 1/\kappa$.

6. Yang-Baxter Equation

The one-loop spin chain of the $\mathbb{Z}_2$ orbifold of $\mathcal{N} = 4$ SYM is known to be integrable [12, 13]. A natural question is whether integrability persists for the $\check{g} \neq g$. We can explore the integrability of the spin chain for the interpolating SCFT by checking the Yang-Baxter equation for the two-body S-matrix. Integrability of the spin chain amounts to the existence of higher conserved quantities beyond the momentum and the Hamiltonian, which would imply exact factorization of many-body scattering into a sequence of two-body scatterings. For this to happen it is necessary that different ways to factorize three-body scattering into two-body scatterings should commute: the Yang-Baxter equation expresses this consistency condition.

The two-body S-matrix of our theory factorizes into the S-matrix for the $SU(2)_L$ sector times the S-matrix for the $SU(2)_R$ sector. The Yang-Baxter equation must be satisfied sep-




Figure 3: Yang-Baxter equation in each $SU(2)$ sector. Simple lines represent Q impurities, double lines $\bar{Q}$ impurities.

arately in each sector. Clearly this is a sufficient condition for the full Yang-Baxter equation to hold; it is also a necessary condition since we can always restrict the asymptotic states to one sector by setting their quantum numbers in the other sector to be highest weights. In each sector, the Yang-Baxter equation is represented by the diagram of figure 3, and reads explicitly

$$S_{IJ}^{\mathcal{MN}}(p_1, p_2) \check{S}_{NK}^{\mathcal{LK}'}(p_1, p_3) S_{ML}^{\mathcal{I}'\mathcal{J}'}(p_2, p_3) = \check{S}_{LP}^{\mathcal{J}'\mathcal{K}'}(p_1, p_2) S_{LN}^{\mathcal{I}'\mathcal{L}}(p_1, p_3) \check{S}_{JK}^{\mathcal{NP}}(p_2, p_3) \quad (6.1)$$

Using the decomposition introduced in the previous section, we can write the left-hand side as

$$\begin{aligned} & S_{IJ}^{\mathcal{MN}}(p_1, p_2) \check{S}_{NK}^{\mathcal{LK}'}(p_1, p_3) S_{ML}^{\mathcal{I}'\mathcal{J}'}(p_2, p_3) \\ &= A\check{A}\check{A}\delta_{\mathcal{K}}^{\mathcal{K}'}\delta_{\mathcal{I}}^{\mathcal{I}'}\delta_{\mathcal{J}}^{\mathcal{J}'} + A\check{B}\check{B}g_{\mathcal{JK}}\delta_{\mathcal{I}}^{\mathcal{K}'}g^{\mathcal{I}'\mathcal{J}'} + B\check{B}Ag_{\mathcal{IJ}}\delta_{\mathcal{K}}^{\mathcal{I}'}g^{\mathcal{J}'\mathcal{K}'} \\ &+ (A\check{A}\check{B} + B\check{A}\check{A} + 2B\check{A}\check{B} + B\check{B}\check{B})\delta_{\mathcal{K}}^{\mathcal{K}'}g_{\mathcal{IJ}}g^{\mathcal{I}'\mathcal{J}'} + A\check{B}Ag_{\mathcal{JK}}g^{\mathcal{J}'\mathcal{K}'}\delta_{\mathcal{I}}^{\mathcal{I}'} \end{aligned}$$

We have suppressed the momentum arguments with the convention that the first symbol in each term is a function of (p_1, p_2) , the second is function of (p_1, p_3) and the third (p_2, p_3) . Similarly, for the right-hand side

$$\begin{aligned} & \check{S}_{LP}^{\mathcal{J}'\mathcal{K}'}(p_1, p_2) S_{LN}^{\mathcal{I}'\mathcal{L}}(p_1, p_3) \check{S}_{JK}^{\mathcal{NP}}(p_2, p_3) \\ &= \check{A}\check{A}\check{A}\delta_{\mathcal{I}}^{\mathcal{I}'}\delta_{\mathcal{J}}^{\mathcal{J}'}\delta_{\mathcal{K}}^{\mathcal{K}'} + \check{A}\check{B}\check{B}g^{\mathcal{I}'\mathcal{J}'}g_{\mathcal{JK}}\delta_{\mathcal{I}}^{\mathcal{K}'} + \check{B}\check{B}Ag^{\mathcal{J}'\mathcal{K}'}g_{\mathcal{IJ}}\delta_{\mathcal{K}}^{\mathcal{I}'} \\ &+ \check{A}\check{B}\check{A}g_{\mathcal{IJ}}g^{\mathcal{I}'\mathcal{J}'}\delta_{\mathcal{K}}^{\mathcal{K}'} + (\check{A}\check{A}\check{B} + \check{B}\check{A}\check{A} + 2\check{B}\check{A}\check{B} + \check{B}\check{B}\check{B})g^{\mathcal{J}'\mathcal{K}'}\delta_{\mathcal{I}}^{\mathcal{I}'}g_{\mathcal{JK}} \end{aligned}$$

Collecting the terms with the same index structure, the Yang-Baxter equation in each $SU(2)$




Figure 4: Example of $SU(2)$ index flow. Collection of the terms with index structure $g_{\mathcal{I}\mathcal{J}}\delta_{\mathcal{K}}^{\mathcal{K}'}g^{\mathcal{I}'\mathcal{J}'}$ gives rise to the 4th YB equation. Similarly, other Yang-Baxter equations also can be understood in graphical manner.

sector reduces to the following five equations:

$$A\check{A}A = \check{A}A\check{A} \quad (6.2)$$

$$A\check{B}B = \check{A}B\check{B} \quad (6.3)$$

$$B\check{B}A = \check{B}B\check{A} \quad (6.4)$$

$$2B\check{A}B + A\check{A}B + B\check{A}A + B\check{B}B = \check{A}B\check{A} \quad (6.5)$$

$$A\check{B}A = 2\check{B}A\check{B} + \check{A}A\check{B} + \check{B}A\check{A} + \check{B}B\check{B}. \quad (6.6)$$

At the orbifold point, $\kappa = 1/\kappa = 1$ and thus $A = \check{A}$, $B = \check{B}$: the first three equations are trivial; the forth and fifth become equivalent. In both the $SU(2)_L$ and $SU(2)_R$ sectors (which are in fact equivalent for $\kappa = 1$), the remaining equation is easily checked. Thus as expected, the Yang-Baxter equation is satisfied at the orbifold point. We then find that YB is *violated* as we move away from the orbifold point, for all $\kappa \in (0, 1)$, showing conclusively that the spin chain of the interpolating theory is *not* integrable for general κ . To our surprise however, YB holds again in the SCQCD limit $\kappa \rightarrow 0$! We take this as a hint that planar $\mathcal{N} = 2$ SCQCD might be integrable, at least at one loop.

7. Discussion

Ordinarily, verification of the Yang-Baxter equation for the two-magnon S-matrix counts as strong evidence for integrability. In our case, however, for κ *strictly* zero, the elementary Q impurities “freeze”, and only $Q\bar{Q}$ dimers can propagate on the chain. Correspondingly, the Q dispersion relation becomes momentum-independent,

$$E_Q(p; \kappa) = 2(1 - \kappa)^2 + 8\kappa \sin^2\left(\frac{p}{2}\right) \xrightarrow{\kappa \rightarrow 0} 2, \quad (7.1)$$

and the S-matrix also degenerates to a simple expression. Verification of YB strictly at $\kappa = 0$ may then appear like an accident due to this degenerate limit. What we find more significant, and non-trivial evidence for integrability, is that the integrable point $\kappa = 0$ is reached smoothly, with YB failing infinitesimally for infinitesimal κ – this is clear since the S-matrices are analytic (rational) functions of κ . This smooth behavior is illustrated in figure




Figure 5: The figure shows the differences between the left and right-hand sides of the five Yang-Baxter equations, as a function of κ , for the specific choice of momenta $p_1 = 0.3$, $p_2 = 0.8$ and $p_3 = 1.4$. The blue, red, green, orange and purple curves show $(l.h.s)-(r.h.s)$ for the first to fifth equation.

5, where we plot the differences between the left and right hand sides of the five equations (6.2–6.6) (for some choice of the momenta).

An elegant way to conclusively prove one-loop integrability at $\kappa = 0$ would be to exhibit the algebraic Bethe ansatz for the SCQCD spin chain. The simplest guess for the R-matrix does not appear to work [32], but the issue deserves further investigation.

Another possible approach is to *assume* integrability to derive Bethe equations for the periodic chain, and then check whether their (numerical) solutions agree with the solutions obtained by direct diagonalization of the Hamiltonian. This is not entirely straightforward, because we cannot work strictly at $\kappa = 0$. The naive Bethe equations at $\kappa = 0$ have no interesting solutions for finite values of the Bethe roots – the non-trivial dynamics is hidden in Bethe roots with infinite imaginary parts (in the momentum variable). We saw this phenomenon in the evolution of the bound states as $\kappa \rightarrow 0$, where the individual magnon momenta behave as $i \log \kappa$. Taking the SCQCD limit $\kappa \rightarrow 0$ too early we lose information about the bound states. (It is conceivable that the failure of the (simplest) algebraic Bethe ansatz is also due to this order-of-limits issue.) Nevertheless, it makes sense to write Bethe equations for small κ , viewed as a regulator to be removed at the end of the calculation. We can also calculate the S-matrix of the bound states, by using the fusion procedure for infinitesimal κ , and check *their* YB equation in the SCQCD limit. The consistency of this approach should follow from

the smoothness of the $\kappa \rightarrow 0$ limit.

A natural extension of our work is the calculation of one-loop dilation operator in the complete theory, including fermions and derivatives [33]. Let us briefly comment on the symmetry structure of the complete spin chain. As is well-known, the symmetry of the $\mathcal{N} = 4$ spin chain in the excitation picture is $PSU(2|2)_L \times PSU(2|2)_R \times \mathbb{R}$, where the central factor $\mathbb{R}$ is identified with the Hamiltonian. The $\mathbb{Z}_2$ orbifold projection preserves the $PSU(2|2)_R$ in the “right” sector (this is a subgroup of the $\mathcal{N} = 2$ superconformal group $SU(2, 2|2)$), but breaks $PSU(2|2)_L$ to the bosonic subgroup $SU(2)_L \times SU(2)_\alpha$, where $SU(2)_\alpha$ denotes the left-handed Lorentz symmetry. At the orbifold point $\kappa = 1$, the breaking is only due to a global twist of the chain, while locally the symmetry is the same as in $\mathcal{N} = 4$, but for $\kappa \neq 1$ the symmetry is truly broken. All in all, the symmetry of the spin chain of the interpolating theory is $SU(2)_L \times SU(2)_\alpha \times PSU(2|2)_R \times \mathbb{R}$. In this paper we have found that in the two-body S-matrix of Q impurities has a left $\times$ right factorization, and we expect this feature to persist for the full chain.

An obvious question is whether symmetry is sufficient to fix the form of the S-matrix, as it does to all loops in $\mathcal{N} = 4$ SYM (up to an overall scalar factor). While unlikely for S_L , this is likely for S_R , which has a large supergroup symmetry. In fact, the symmetry in the right sector of the interpolating SCFT the same as in (either sector of) $\mathcal{N} = 4$ SYM. The S_R matrix of $\mathcal{N} = 4$ is uniquely fixed, up to an overall scalar factor, from the (centrally extended) $SU(2|2)_R$ symmetry [34]. But our results for S_R in the interpolating theory are definitely different (for $\kappa \neq 1$) from the $\mathcal{N} = 4$ results. This is clear already in the scalar sector studied in this paper, by inspection of the S-matrix of the $Q_{\mathcal{I}\hat{+}}$ impurities. This discrepancy is explained by the fact in our case the magnons transform in a reducible representation of $SU(2|2)$ (two copies of the fundamental representation). It will be interesting to see whether these assumptions can be relaxed to reproduce (and possibly uniquely fix) Remarkably, it is still possible to use symmetry to fix uniquely the form of the S_R matrix in the interpolating theory, up to a free parameter that can be identified with κ . This analysis will be presented elsewhere [35].

Finally it would be very interesting to evaluate the two-body S-matrix at strong coupling, in the dual string sigma-model, and see whether it has the same κ dependence as the perturbative S-matrix. Failure of integrability for generic κ is not an issue here, since we would not be using in any way factorization of n -body scattering, but rather focus on the two-body S-matrix, which we expect to have a smooth interpolation from weak to strong coupling. The sigma-model at the orbifold point is well-known, and moving away from the orbifold point corresponds to changing the value of a theta angle β (the period of the NSNS B -field through the collapsed cycle of the orbifold) [36, 37]. The orbifold point corresponds to $\beta = 1/2$, while the SCQCD limit corresponds to $\beta \rightarrow 0$. From the dual side, it is natural to expect integrability precisely at the two extrema 0 and $1/2$, but not for generic values of the B -field. A toy model for this behavior is the $O(3)$ sigma-model in a magnetic field [38].

One of our original motivations was to collect “bottom-up” clues about the string dual of $\mathcal{N} = 2$ SCQCD. While firm conclusions will have to wait a higher-order (all order?) analysis,

we can already see a qualitative agreement with the “top-down” approach of our previous paper [1]. We argued that $\mathcal{N} = 2$ SCQCD is dual to a non-critical string background, with seven geometric dimensions, containing both an AdS_5 and an S^1 factor. Rotation in S^1 corresponds to the $U(1)_r$ quantum number. In lightcone quantization of the sigma-model, the lightcone coordinates would be obtained by combining this S^1 and the timelike direction of AdS_5 . We then expect five bosonic gapless excitations, four associated to the transverse AdS coordinates and one to the seventh dimension. The vacuum of the lightcone sigma-model corresponds to chiral vacuum $\text{Tr } \phi^\ell$ of the spin chain, while the four AdS excitations correspond to derivative impurities on the chain. In the scalar sector that we have studied in this paper, *one* gapless excitation is then expected, the one corresponding to the seventh dimension: just what we found, the gapless magnon $T(p)$. As $\kappa \rightarrow 0$, the Q impurities, carriers of the $SU(2)_L \times SU(2)_R$ quantum numbers associated with the three extra dimensions (the transverse S^3 , see [1] for details), become non-dynamical, and only their composite bound state $T(p)$ survives as a gapless mode. We interpret this phenomenon as the field theory counterpart of the transition from the critical to the non-critical background.

Acknowledgements

It is a pleasure thank Vladimir Korepin, Pedro Liendo, Juan Maldacena, Joe Minahan, Joe Polchinski, Soo-Jong Rey, Martin Rocek, Radu Roiban, Matthias Staudacher, Pedro Vieira, and Kostya Zarembo for useful discussions. This work was supported in part by DOE grant DEFG-0292-ER40697 and by NSF grant PHY-0653351-001. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

A. Simplified computation of the one-loop dilation operator

In this appendix we determine the one-loop spin-chain Hamiltonian by a simple shortcut. The interactions contributing to $H_{k,k+1}$ at one loop are listed schematically in figure 1. The first and second interactions (self-energy and gluon exchange) in figure 1 are proportional to the identity operator in $V_k \otimes V_{k+1}$, while the non-trivial tensorial structures are contributed only by the third diagram (quartic interaction). The idea is to evaluate explicitly the third diagram, and to fix the terms proportional to the identity by requiring that the anomalous dimensions of a few protected operators vanish.

A.1 SCQCD

Let us recall our notations. The indices $\mathbf{p}, \mathbf{q} = \pm$ label the $U(1)_r$ charges of ϕ and $\bar{\phi}$, in other terms we define $\phi^- \equiv \phi$, $\phi^+ \equiv \bar{\phi}$, and $g_{\mathbf{p}\mathbf{q}} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. The elements of the Hamiltonian due to




Figure 6: The color/ flavor structure of the quartic vertex. The solid black line represents the flow of the color index while the dotted blue line show the flow of the flavor index. Diagram (a) shows the ϕ^4 interaction vertex, whose contribution is proportional to N_c as compared to the tree level. In (b) the $Q^2\phi^2$ interaction vertex has a factor of N_f/N_c compared to (a) because of the presence of one flavor loop. The Q^4 vertex in (c) has an additional factor of $(N_f/N_c)^2$ compared to (a) due to the presence of two flavor loops. Diagram (d), however, does not carry any additional N_f/N_c factors.

quartic vertices are:

$$\langle \phi_{\mathbf{p}'} \phi_{\mathbf{q}'} | H | \phi^{\mathbf{p}} \phi^{\mathbf{q}} \rangle_{\phi^4} = \delta_{\mathbf{p}'}^{\mathbf{p}} \delta_{\mathbf{q}'}^{\mathbf{q}} + g^{\mathbf{p}\mathbf{q}} g_{\mathbf{p}'\mathbf{q}'} - 2\delta_{\mathbf{q}'}^{\mathbf{p}} \delta_{\mathbf{p}'}^{\mathbf{q}} \quad (\text{A.1})$$

$$\langle \phi_{\mathbf{p}'} \phi_{\mathbf{q}'} | H | Q_{\mathcal{I}} \bar{Q}^{\mathcal{J}} \rangle_{Q^2\phi^2} = \sqrt{\frac{N_f}{N_c}} g_{\mathbf{p}'\mathbf{q}'} \delta_{\mathcal{I}}^{\mathcal{J}} \quad (\text{A.2})$$

$$\langle \bar{Q}^{\mathcal{I}'} Q_{\mathcal{J}'} | H | Q_{\mathcal{I}} \bar{Q}^{\mathcal{J}} \rangle_{Q^4} = \frac{N_f}{N_c} (2\delta_{\mathcal{I}}^{\mathcal{I}'} \delta_{\mathcal{J}'}^{\mathcal{J}} - \delta_{\mathcal{I}}^{\mathcal{J}} \delta_{\mathcal{J}'}^{\mathcal{I}'}) \quad (\text{A.3})$$

$$\langle Q_{\mathcal{J}'} \bar{Q}^{\mathcal{I}'} | H | \bar{Q}^{\mathcal{J}} Q_{\mathcal{I}} \rangle_{Q^4} = 2\delta_{\mathcal{I}}^{\mathcal{J}} \delta_{\mathcal{J}'}^{\mathcal{I}'} - \delta_{\mathcal{I}}^{\mathcal{I}'} \delta_{\mathcal{J}'}^{\mathcal{J}} \quad (\text{A.4})$$

The factors of $\frac{N_f}{N_c}$ are explained in figure 6. Figures 6a,6b,6c,6d correspond to equations (A.1,A.2,A.3,A.4) respectively. This fixes the Hamiltonian up to the terms proportional to the identity,

$$H_{k,k+1} = \begin{matrix} & \phi^{\mathbf{p}} \phi^{\mathbf{q}} & Q_{\mathcal{I}} \bar{Q}^{\mathcal{J}} & \bar{Q}^{\mathcal{K}} Q_{\mathcal{L}} & Q_{\mathcal{I}} \phi^{\mathbf{p}} \\ \begin{matrix} \phi_{\mathbf{p}'} \phi_{\mathbf{q}'} \\ \bar{Q}^{\mathcal{I}'} Q_{\mathcal{J}'} \\ Q_{\mathcal{K}'} \bar{Q}^{\mathcal{L}'} \\ \bar{Q}^{\mathcal{I}'} \phi_{\mathbf{p}'} \end{matrix} & \begin{pmatrix} \alpha \delta_{\mathbf{p}'}^{\mathbf{p}} \delta_{\mathbf{q}'}^{\mathbf{q}} + g^{\mathbf{p}\mathbf{q}} g_{\mathbf{p}'\mathbf{q}'} - 2\delta_{\mathbf{q}'}^{\mathbf{p}} \delta_{\mathbf{p}'}^{\mathbf{q}} & \sqrt{\frac{N_f}{N_c}} g_{\mathbf{p}'\mathbf{q}'} \delta_{\mathcal{I}}^{\mathcal{J}} & 0 & 0 \\ \sqrt{\frac{N_f}{N_c}} g^{\mathbf{p}\mathbf{q}} \delta_{\mathcal{J}'}^{\mathcal{I}'} & \beta \delta_{\mathcal{I}}^{\mathcal{I}'} \delta_{\mathcal{J}'}^{\mathcal{J}} - \delta_{\mathcal{I}}^{\mathcal{J}} \delta_{\mathcal{J}'}^{\mathcal{I}'} \frac{N_f}{N_c} & 0 & 0 \\ 0 & 0 & \gamma \delta_{\mathcal{K}'}^{\mathcal{K}} \delta_{\mathcal{L}'}^{\mathcal{L}} + 2\delta_{\mathcal{L}}^{\mathcal{K}} \delta_{\mathcal{K}'}^{\mathcal{L}} & 0 \\ 0 & 0 & 0 & \eta \delta_{\mathcal{I}}^{\mathcal{I}'} \delta_{\mathbf{p}'}^{\mathbf{p}} \end{pmatrix} \end{matrix}$$

We can now find the coefficients α, β, γ and η from knowledge of the protected spectrum. Vanishing of the anomalous dimension of $\text{Tr} \phi^k$ gives $\alpha = 2$. Another protected multiplet is the multiplet containing the stress-energy tensor. Its superconformal primary, called $\text{Tr } T$,

has $R, r = 0$ and $\Delta = 2$. Hence, it is a linear combination of $\text{Tr}[Q_{\mathcal{I}}\bar{Q}^{\mathcal{I}}]$ and $\text{Tr}[\phi\bar{\phi}]$. The restriction of the Hamiltonian to this subspace is

$$H = \begin{pmatrix} \text{Tr}[\phi\bar{\phi}] & \text{Tr}[\mathcal{M}_1] \\ \text{Tr}[\phi\bar{\phi}] & 2\sqrt{\frac{2N_f}{N}} \\ \text{Tr}[\mathcal{M}_1] & (\beta + \gamma) - 2(\frac{N_f}{N_c} - 2) \end{pmatrix} \quad (\text{A.5})$$

This matrix must have a zero at the superconformal point $N_f = 2N_c$, yielding $\beta + \gamma = 4$. Finally, the fact that $\text{Tr} T\phi$ is also a protected operator gives the relation $\beta + 2\eta = 8$. We started with four coefficients $\alpha, \beta, \gamma, \eta$ and imposed three relations. The undetermined degrees of freedom corresponds to the “gauge” freedom of adding to the nearest neighbor Hamiltonian terms that vanish upon evaluating the full H on a closed chain. We may solve the constraints by writing

$$\alpha = 2, \quad \beta = 4 + \frac{1}{2}(1 + \xi), \quad \gamma = -\frac{1}{2}(1 + \xi), \quad \eta = \frac{1}{4}(7 - \xi), \quad (\text{A.6})$$

where ξ is the arbitrary gauge parameter. The resulting Hamiltonian is in perfect agreement (for $N_f = 2N_c$) with the answer (3.3) obtained by the slightly lengthier route of explicit evaluating all relevant one-loop diagrams. All in all, this confirms our understanding of the protected spectrum.

A.2 Interpolating SCFT

We can repeat the same exercise for the interpolating SCFT. The quartic vertices give

$$\langle \phi_{\mathbf{p}'}\phi_{\mathbf{q}'}|\phi^{\mathbf{p}}\phi^{\mathbf{q}}\rangle_{\phi^4} = \delta_{\mathbf{p}'}^{\mathbf{p}}\delta_{\mathbf{q}'}^{\mathbf{q}} + g^{\mathbf{p}\mathbf{q}}g_{\mathbf{p}'\mathbf{q}'} - 2\delta_{\mathbf{q}'}^{\mathbf{p}}\delta_{\mathbf{p}'}^{\mathbf{q}} \quad (\text{A.7})$$

$$\langle \check{\phi}_{\mathbf{p}'}\check{\phi}_{\mathbf{q}'}|\check{\phi}^{\mathbf{p}}\check{\phi}^{\mathbf{q}}\rangle_{\check{\phi}^4} = \kappa^2(\delta_{\mathbf{p}'}^{\mathbf{p}}\delta_{\mathbf{q}'}^{\mathbf{q}} + g^{\mathbf{p}\mathbf{q}}g_{\mathbf{p}'\mathbf{q}'} - 2\delta_{\mathbf{q}'}^{\mathbf{p}}\delta_{\mathbf{p}'}^{\mathbf{q}}) \quad (\text{A.8})$$

$$\begin{aligned} \langle \bar{Q}^{\hat{\mathcal{L}}\mathcal{L}}Q_{\mathcal{K}\hat{\mathcal{K}}}|Q_{\mathcal{I}\hat{\mathcal{I}}}\bar{Q}^{\hat{\mathcal{J}}\mathcal{J}}\rangle_{Q^4} &= 2\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}}\delta_{\mathcal{K}}^{\mathcal{J}}\delta_{\hat{\mathcal{K}}}^{\mathcal{L}}\delta_{\mathcal{I}}^{\mathcal{L}} - \delta_{\mathcal{I}}^{\mathcal{J}}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}}\delta_{\mathcal{K}}^{\mathcal{L}}\delta_{\hat{\mathcal{K}}}^{\mathcal{L}} \\ &+ \kappa^2(2\delta_{\hat{\mathcal{K}}}^{\hat{\mathcal{J}}}\delta_{\mathcal{I}}^{\mathcal{J}}\delta_{\hat{\mathcal{I}}}^{\mathcal{L}}\delta_{\mathcal{K}}^{\mathcal{L}} - \delta_{\mathcal{I}}^{\mathcal{L}}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}}\delta_{\mathcal{K}}^{\mathcal{J}}\delta_{\hat{\mathcal{K}}}^{\hat{\mathcal{J}}}) \end{aligned} \quad (\text{A.9})$$

$$\begin{aligned} \langle Q_{\mathcal{I}\hat{\mathcal{I}}}\bar{Q}^{\hat{\mathcal{J}}\mathcal{J}}|\bar{Q}^{\hat{\mathcal{L}}\mathcal{L}}Q_{\mathcal{K}\hat{\mathcal{K}}}\rangle_{Q^4} &= 2\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}}\delta_{\mathcal{K}}^{\mathcal{J}}\delta_{\hat{\mathcal{K}}}^{\mathcal{L}}\delta_{\mathcal{I}}^{\mathcal{L}} - \delta_{\mathcal{I}}^{\mathcal{J}}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}}\delta_{\mathcal{K}}^{\mathcal{L}}\delta_{\hat{\mathcal{K}}}^{\mathcal{L}} \\ &+ \kappa^2(2\delta_{\hat{\mathcal{K}}}^{\hat{\mathcal{J}}}\delta_{\mathcal{I}}^{\mathcal{J}}\delta_{\hat{\mathcal{I}}}^{\mathcal{L}}\delta_{\mathcal{K}}^{\mathcal{L}} - \delta_{\mathcal{I}}^{\mathcal{L}}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}}\delta_{\mathcal{K}}^{\mathcal{J}}\delta_{\hat{\mathcal{K}}}^{\hat{\mathcal{J}}}) \end{aligned} \quad (\text{A.10})$$

$$\langle \phi_{\mathbf{p}'}\phi_{\mathbf{q}'}|Q_{\mathcal{I}\hat{\mathcal{I}}}\bar{Q}^{\hat{\mathcal{J}}\mathcal{J}}\rangle_{Q^2\phi^2} = g_{\mathbf{p}'\mathbf{q}'}\delta_{\mathcal{I}}^{\mathcal{J}}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}} \quad (\text{A.11})$$

$$\langle \check{\phi}_{\mathbf{p}'}\check{\phi}_{\mathbf{q}'}|\bar{Q}^{\hat{\mathcal{J}}\mathcal{J}}Q_{\mathcal{I}\hat{\mathcal{I}}}\rangle_{Q^2\check{\phi}^2} = \kappa^2 g_{\mathbf{p}'\mathbf{q}'}\delta_{\mathcal{I}}^{\mathcal{J}}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}} \quad (\text{A.12})$$

$$\langle \bar{Q}^{\hat{\mathcal{J}}\mathcal{J}}\check{\phi}_{\mathbf{q}}|\phi^{\mathbf{p}}Q_{\mathcal{I}\hat{\mathcal{I}}}\rangle_{\phi Q\check{\phi}\bar{Q}} = -2\kappa\delta_{\mathbf{q}}^{\mathbf{p}}\delta_{\mathcal{I}}^{\mathcal{J}}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}} \quad (\text{A.13})$$

$$\langle \phi^{\mathbf{p}}\bar{Q}^{\hat{\mathcal{J}}\mathcal{J}}|Q_{\mathcal{I}\hat{\mathcal{I}}}\check{\phi}_{\mathbf{q}}\rangle_{\phi Q\check{\phi}\bar{Q}} = -2\kappa\delta_{\mathbf{q}}^{\mathbf{p}}\delta_{\mathcal{I}}^{\mathcal{J}}\delta_{\hat{\mathcal{I}}}^{\hat{\mathcal{J}}} \quad (\text{A.14})$$

The first four elements can have additional identity pieces. They are easily determined by imposing the symmetry under $g \leftrightarrow \check{g}$, $Q \leftrightarrow \bar{Q}$ and $\phi \leftrightarrow \check{\phi}$ and by requiring the Hamiltonian

to reduce to that of SCQCD in the limit $\kappa \rightarrow 0$. The one loop Hamiltonian (3.16) is precisely reproduced by this method.

B. The Hamiltonian for SCQCD in the Dimer Picture

In this appendix we rewrite the Hamiltonian for SCQCD as acting on adjoint fields and dimers $Q_{\mathcal{I}}\bar{Q}^{\mathcal{J}}$, regarded as basic objects. We define the singlet combination $\mathcal{M} = \frac{1}{\sqrt{2}}\mathcal{M}_{\mathcal{I}}^{\mathcal{J}}\delta_{\mathcal{J}}^{\mathcal{I}}$ and the triplet $\mathcal{M}^i = \frac{1}{\sqrt{2}}\mathcal{M}_{\mathcal{I}}^{\mathcal{J}}(\sigma^i)_{\mathcal{J}}^{\mathcal{I}}$, where σ^i are three Pauli matrices. These can be rewritten in an $SO(4)$ notation as $\mathcal{M}^m = \frac{1}{\sqrt{2}}\mathcal{M}_{\mathcal{I}}^{\mathcal{J}}(\sigma^m)_{\mathcal{J}}^{\mathcal{I}}$, where $m = 0, \dots, 3$ and $\sigma^0 \equiv \mathbb{I}_{2 \times 2}$.

Consider the action of H on following sequence in the spin chain,

$$\begin{array}{ccccccc}
 \phi^{\mathfrak{p}} & & Q_{\mathcal{I}} & & \bar{Q}^{\mathcal{J}} & & \phi^{\mathfrak{q}} \\
 & \frac{1}{2}(3 + \frac{\xi}{2}) & & (5 - \frac{\xi}{2})\mathbb{I}_{QQ} - 2\mathbb{K}_{QQ} & & \frac{1}{2}(3 + \frac{\xi}{2}) & \\
 & \downarrow & & \downarrow & & \downarrow & \\
 \phi_{\mathfrak{p}'} & & \bar{Q}^{\mathcal{I}'} & & Q_{\mathcal{J}'} & & \phi_{\mathfrak{q}'}
 \end{array} \tag{B.1}$$

In the new picture, where $\mathcal{M}$ is regarded as a basic impurity, the middle term $(5 - \frac{\xi}{2})\mathbb{I}_{QQ} - 2\mathbb{K}_{QQ}$ is the “self energy” of $\mathcal{M}$, and we split it evenly between the $\phi\mathcal{M}$ and $\mathcal{M}\phi$ matrix elements. So we write

$$\begin{aligned}
 \langle \dots \phi_{\mathfrak{p}'} \bar{\mathcal{M}}^{\mathcal{I}'} \dots | H | \dots \phi^{\mathfrak{p}} \mathcal{M}_{\mathcal{I}}^{\mathcal{J}} \dots \rangle &= [\frac{1}{2}(3 + \frac{\xi}{2}) + \frac{1}{2}(5 - \frac{\xi}{2})] \delta_{\mathfrak{p}'}^{\mathfrak{p}} \delta_{\mathcal{I}}^{\mathcal{I}'} \delta_{\mathcal{J}}^{\mathcal{J}'} - \delta_{\mathfrak{p}'}^{\mathfrak{p}} \delta_{\mathcal{I}}^{\mathcal{J}} \delta_{\mathcal{J}'}^{\mathcal{I}'} \\
 &= (4\delta_{\mathcal{I}}^{\mathcal{I}'} \delta_{\mathcal{J}'}^{\mathcal{J}} - \delta_{\mathcal{I}}^{\mathcal{J}} \delta_{\mathcal{J}'}^{\mathcal{I}'}) \delta_{\mathfrak{p}'}^{\mathfrak{p}} \\
 \langle \dots \phi_{\mathfrak{p}'} \bar{\mathcal{M}}^{m'} \dots | H | \dots \phi^{\mathfrak{p}} \mathcal{M}^m \dots \rangle &= \delta_{\mathfrak{p}'}^{\mathfrak{p}} \delta^{mm'} (4 - 2\delta^{m0}).
 \end{aligned}$$

Similarly, to find the action of H on two neighboring $\mathcal{M}$ s, we consider the sequence

$$\begin{array}{ccccccc}
 Q_{\mathcal{I}} & & \bar{Q}^{\mathcal{J}} & & Q_{\mathcal{K}} & & \bar{Q}^{\mathcal{L}} \\
 (5 - \frac{\xi}{2})\mathbb{I}_{QQ} - 2\mathbb{K}_{QQ} & & (\frac{\xi}{2} - 1)\mathbb{I}_{QQ} + 2\mathbb{K}_{QQ} & & (5 - \frac{\xi}{2})\mathbb{I}_{QQ} - 2\mathbb{K}_{QQ} & & \\
 \downarrow & & \downarrow & & \downarrow & & \\
 \bar{Q}^{\mathcal{I}'} & & Q_{\mathcal{J}'} & & \bar{Q}^{\mathcal{K}'} & & Q_{\mathcal{L}'}
 \end{array} \tag{B.2}$$

This gives

$$\begin{aligned}
 \langle \dots \bar{\mathcal{M}}^{m'} \bar{\mathcal{M}}^{n'} \dots | H | \dots \mathcal{M}^m \mathcal{M}^n \dots \rangle &= \delta^{mm'} \delta^{nn'} (13 - 4\delta^{m0} - 4\delta^{n0}) \\
 &\quad + \delta^{mn} \delta^{m'n'} - \delta^{mn'} \delta^{nm'} + i\epsilon^{mnn'm'} .
 \end{aligned}$$

References

- [1] A. Gadde, E. Pomoni, and L. Rastelli, *The Veneziano Limit of $\mathcal{N} = 2$ Superconformal QCD: Towards the String Dual of $\mathcal{N} = 2$ $SU(N_c)$ SYM with $N_f = 2N_c$* , [arXiv:0912.4918](#).
- [2] J. A. Minahan and K. Zarembo, *The Bethe-ansatz for $N = 4$ super Yang-Mills*, *JHEP* **03** (2003) 013, [[hep-th/0212208](#)].

- [3] N. Beisert and M. Staudacher, *The $N=4$ SYM Integrable Super Spin Chain*, *Nucl. Phys.* **B670** (2003) 439–463, [[hep-th/0307042](#)].
- [4] M. Staudacher, *The factorized S -matrix of CFT/AdS*, *JHEP* **05** (2005) 054, [[hep-th/0412188](#)].
- [5] N. Beisert and M. Staudacher, *Long-range PSU(2,2|4) Bethe ansatz for gauge theory and strings*, *Nucl. Phys.* **B727** (2005) 1–62, [[hep-th/0504190](#)].
- [6] N. Beisert, B. Eden, and M. Staudacher, *Transcendentality and crossing*, *J. Stat. Mech.* **0701** (2007) P021, [[hep-th/0610251](#)].
- [7] N. Gromov, V. Kazakov, and P. Vieira, *Exact Spectrum of Anomalous Dimensions of Planar $N=4$ Supersymmetric Yang-Mills Theory*, *Phys. Rev. Lett.* **103** (2009) 131601, [[arXiv:0901.3753](#)].
- [8] S.-J. Rey and T. Suyama, *Exact Results and Holography of Wilson Loops in $N=2$ Superconformal (Quiver) Gauge Theories*, [arXiv:1001.0016](#).
- [9] G. Veneziano, *Some Aspects of a Unified Approach to Gauge, Dual and Gribov Theories*, *Nucl. Phys.* **B117** (1976) 519–545.
- [10] G. 't Hooft, *A planar diagram theory for strong interactions*, *Nucl. Phys.* **B72** (1974) 461.
- [11] J. Kinney, J. M. Maldacena, S. Minwalla, and S. Raju, *An index for 4 dimensional super conformal theories*, *Commun. Math. Phys.* **275** (2007) 209–254, [[hep-th/0510251](#)].
- [12] N. Beisert and R. Roiban, *The Bethe ansatz for $Z(S)$ orbifolds of $N = 4$ super Yang- Mills theory*, *JHEP* **11** (2005) 037, [[hep-th/0510209](#)].
- [13] A. Solov'yov, *Bethe Ansatz Equations for General Orbifolds of $N=4$ SYM*, *JHEP* **04** (2008) 013, [[arXiv:0711.1697](#)].
- [14] M. R. Douglas and G. W. Moore, *D-branes, Quivers, and ALE Instantons*, [hep-th/9603167](#).
- [15] A. Dymarsky, I. R. Klebanov, and R. Roiban, *Perturbative gauge theory and closed string tachyons*, *JHEP* **11** (2005) 038, [[hep-th/0509132](#)].
- [16] J. A. Minahan and K. Zarembo, *The Bethe ansatz for superconformal Chern-Simons*, *JHEP* **09** (2008) 040, [[arXiv:0806.3951](#)].
- [17] D. Bak and S.-J. Rey, *Integrable Spin Chain in Superconformal Chern-Simons Theory*, *JHEP* **10** (2008) 053, [[arXiv:0807.2063](#)].
- [18] D. Gaiotto, S. Giombi, and X. Yin, *Spin Chains in $N=6$ Superconformal Chern-Simons-Matter Theory*, *JHEP* **04** (2009) 066, [[arXiv:0806.4589](#)].
- [19] D. Bak, D. Gang, and S.-J. Rey, *Integrable Spin Chain of Superconformal $U(M) \times U(N)$ Chern-Simons Theory*, *JHEP* **10** (2008) 038, [[arXiv:0808.0170](#)].
- [20] O. Aharony, O. Bergman, D. L. Jafferis, and J. Maldacena, *$N=6$ superconformal Chern-Simons-matter theories, $M2$ -branes and their gravity duals*, *JHEP* **10** (2008) 091, [[arXiv:0806.1218](#)].
- [21] O. Aharony, O. Bergman, and D. L. Jafferis, *Fractional $M2$ -branes*, *JHEP* **11** (2008) 043, [[arXiv:0807.4924](#)].

- [22] J. A. Minahan, W. Schulgin, and K. Zarembo, *Two loop integrability for Chern-Simons theories with $N=6$ supersymmetry*, *JHEP* **03** (2009) 057, [[arXiv:0901.1142](#)].
- [23] J. Minahan, O. Ohlsson Sax, and C. Sieg, *Magnon dispersion to four loops in the ABJM and ABJ models*, *J.Phys.A* **A43** (2010) 275402, [[arXiv:0908.2463](#)].
- [24] P. Di Vecchia and A. Tanzini, *$N = 2$ super Yang-Mills and the XXZ spin chain*, *J. Geom. Phys.* **54** (2005) 116–130, [[hep-th/0405262](#)].
- [25] V. K. Dobrev and V. B. Petkova, *ON THE GROUP THEORETICAL APPROACH TO EXTENDED CONFORMAL SUPERSYMMETRY: CLASSIFICATION OF MULTIPLETS*, *Lett. Math. Phys.* **9** (1985) 287–298.
- [26] V. K. Dobrev and V. B. Petkova, *GROUP THEORETICAL APPROACH TO EXTENDED CONFORMAL SUPERSYMMETRY: FUNCTION SPACE REALIZATIONS AND INVARIANT DIFFERENTIAL OPERATORS*, *Fortschr. Phys.* **35** (1987) 537.
- [27] V. K. Dobrev and V. B. Petkova, *All Positive Energy Unitary Irreducible Representations of Extended Conformal Supersymmetry*, *Phys. Lett.* **B162** (1985) 127–132.
- [28] F. A. Dolan and H. Osborn, *On short and semi-short representations for four dimensional superconformal symmetry*, *Ann. Phys.* **307** (2003) 41–89, [[hep-th/0209056](#)].
- [29] V. K. Dobrev, *Characters of the positive energy UIRs of $D = 4$ conformal supersymmetry*, *Phys. Part. Nucl.* **38** (2007) 564–609, [[hep-th/0406154](#)].
- [30] S. Gukov, *Comments on $N = 2$ AdS orbifolds*, *Phys. Lett.* **B439** (1998) 23–28, [[hep-th/9806180](#)].
- [31] Y. Nakayama, *Index for orbifold quiver gauge theories*, *Phys. Lett.* **B636** (2006) 132–136, [[hep-th/0512280](#)].
- [32] A. Gadde and W. Yan, *Private Communication*, .
- [33] P. Liendo, E. Pomoni, and L. Rastelli, *The Complete One-Loop Dilation Operator of $N=2$ SuperConformal QCD*, [arXiv:1105.3972](#).
- [34] N. Beisert, *The $su(2/2)$ dynamic S -matrix*, *Adv. Theor. Math. Phys.* **12** (2008) 945, [[hep-th/0511082](#)].
- [35] A. Gadde and L. Rastelli, *Twisted Magnons*, [arXiv:1012.2097](#).
- [36] A. E. Lawrence, N. Nekrasov, and C. Vafa, *On conformal field theories in four dimensions*, *Nucl. Phys.* **B533** (1998) 199–209, [[hep-th/9803015](#)].
- [37] I. R. Klebanov and N. A. Nekrasov, *Gravity duals of fractional branes and logarithmic RG flow*, *Nucl. Phys.* **B574** (2000) 263–274, [[hep-th/9911096](#)].
- [38] A. B. Zamolodchikov and A. B. Zamolodchikov, *Massless factorized scattering and sigma models with topological terms*, *Nucl. Phys.* **B379** (1992) 602–623.